

CORRECTED VERSION

Rec'd PCT/PTO 02 OCT 2005

10/551840

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
14 October 2004 (14.10.2004)

PCT

(10) International Publication Number
WO 2004/087213 A1

(51) International Patent Classification⁷: A61K 47/10,
38/24

(21) International Application Number:
PCT/EP2004/050432

(22) International Filing Date: 2 April 2004 (02.04.2004)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
03100882.4 2 April 2003 (02.04.2003) EP
03101543.1 27 May 2003 (27.05.2003) EP
03101828.6 20 June 2003 (20.06.2003) EP

(71) Applicant (for all designated States except US): ARES
TRADING S.A. [CH/CH]; Zone Industrielle de l'Ouriet-
taz, CH-1170 Aubonne (CH).

(72) Inventors; and

(75) Inventors/Applicants (for US only): SAMARITANI,
Fabrizio [IT/IT]; Via Luigi Chiala 130, I-00139 Roma
(IT). DONATI, Piergiorgio [IT/CH]; Rue de Lausanne
66, CH-1110 Morges (CH).

(74) Agent: HASSA, Juergen; Serson International SA, Intel-
lectual Property, 12, Chemin des Aulx, CH-1228 Plan-les-
Ouates (CH).

(81) Designated States (unless otherwise indicated, for every
kind of national protection available): AE, AG, AL, AM,

AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN,
CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI,
GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE,
KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD,
MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG,
PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM,
TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM,
ZW.

(84) Designated States (unless otherwise indicated, for every
kind of regional protection available): ARIPO (BW, GH,
GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),
Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), Euro-
pean (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR,
GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI, SK,
TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW,
ML, MR, NE, SN, TD, TG).

Published:

- with international search report
- with sequence listing part of description published sepa-
rately in electronic form and available upon request from
the International Bureau

(48) Date of publication of this corrected version:

17 March 2005

(15) Information about Correction:

see PCT Gazette No. 11/2005 of 17 March 2005, Section II

For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.

(54) Title: LIQUID PHARMACEUTICAL FORMULATIONS OF FSH AND LH TOGETHER WITH A NON-IONIC SURFAC-
TANT

(57) Abstract: The invention relates to the field of pharmaceutical formulations of follicle -stimulating hormone (FSH), luteinising
hormone (LH), and mixtures of FSH and luteinising hormone (LH), and to methods of producing such formulations. The invention
provides a liquid or freeze-dried formulation of FSH, or LH, or FSH and LH comprising a surfactant selected from Pluronic[®] F77,
Pluronic F87, Pluronic F88 and Pluronic F68.

WO 2004/087213 A1

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
14 October 2004 (14.10.2004)

PCT

Rec'd PCT/PTO 02 OCT 2005



10/551840

(10) International Publication Number
WO 2004/087213 A1

- (51) International Patent Classification⁷: A61K 47/10, 38/24
- (21) International Application Number: PCT/EP2004/050432
- (22) International Filing Date: 2 April 2004 (02.04.2004)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:
- | | | |
|------------|---------------------------|----|
| 03100882.4 | 2 April 2003 (02.04.2003) | EP |
| 03101543.1 | 27 May 2003 (27.05.2003) | EP |
| 03101828.6 | 20 June 2003 (20.06.2003) | EP |
- (71) Applicant (for all designated States except US): ARES TRADING S.A. [CH/CH]; Zone Industrielle de l'Ourietaz, CH-1170 Aubonne (CH).
- (72) Inventors; and
- (75) Inventors/Applicants (for US only): SAMARITANI, Fabrizio [IT/IT]; Via Luigi Chiala 130, I-00139 Roma (IT). DONATI, Piergiorgio [IT/CH]; Rue de Lausanne 66, CH-1110 Morges (CH).
- (74) Agent: HASSA, Juergen; Serono International SA, Intellectual Property, 12, Chemin des Aulx, CH-1228 Plan-les-Ouates (CH).
- (81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.
- (84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).
- Published:
- with international search report
 - before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: LIQUID PHARMACEUTICAL FORMULATIONS OF FSH AND LH TOGETHER WITH A NON-IONIC SURFACTANT

(57) Abstract: The invention relates to the field of pharmaceutical formulations of follicle -stimulating hormone (FSH), luteinising hormone (LH), and mixtures of FSH and luteinising hormone (LH), and to methods of producing such formulations. The invention provides a liquid or freeze-dried formulation of FSH, or LH, or FSH and LH comprising a surfactant selected from Pluronic[®] F77, Pluronic F87, Pluronic F88 and Pluronic F68.

WO 2004/087213 A1

LIQUID PHARMACEUTICAL FORMULATIONS OF FSH AND LH TOGETHER WITH A NON-IONIC SURFACTANT

Field of Invention

The invention relates to the field of pharmaceutical formulations of follicle-stimulating hormone (FSH), formulations of luteinising hormone (LH), and mixtures of FSH and luteinising hormone (LH); as well as to methods of producing such formulations.

Background of the invention

Follicle-stimulating hormone (FSH), luteinising hormone (LH) and chorionic gonadotrophin (CG) are injectable proteins falling into the class of gonadotrophins. FSH, LH and hCG are used alone and in combination in the treatment of infertility and reproductive disorders in both female and male patients.

In nature, FSH and LH are produced by the pituitary gland. For pharmaceutical use, FSH and LH and their variants may be produced recombinantly (rFSH and rLH), or they may be produced from the urine of postmenopausal women (uFSH and uLH).

FSH is used in female patients in ovulation induction (OI) and in controlled ovarian hyperstimulation (COH) for assisted reproductive technologies (ART). In a typical treatment regimen for ovulation induction, a patient is administered daily injections of FSH or a variant (about 75 to 300 IU FSH/day) for a period of from about 6 to about 12 days. In a typical treatment regimen for controlled ovarian hyperstimulation, a patient is administered daily injections of FSH or a variant (about 150-600 IU FSH/day) for a period of from about 6 to about 12 days.

FSH is also used to induce spermatogenesis in men suffering from oligospermia. A regimen using 150 IU FSH 3 times weekly in combination with 2'500 IU hCG twice weekly has been successful in achieving an improvement in sperm count in men suffering from hypogonadotrophic hypogonadism¹.

LH is used in female patients in combination with FSH in OI and in COH, particularly in those patients having very low endogenous LH levels or resistance to LH, such as women suffering from hypogonadotrophic hypogonadism (HH, WHO group I) or older patients (i.e. 35 years or older), and patients in which embryo implantation or early miscarriage is a problem. LH in combination with FSH has traditionally been available in a preparation called human menopausal gonadotrophins (hMG)

extracted from the urine of postmenopausal women. hMG has a 1:1 ratio of FSH:LH activity.

CG acts at the same receptor as LH and elicits the same responses. CG has a longer circulation half-life than LH and is therefore commonly used as a long-acting source of LH-activity. CG is used in OI and COH regimens to mimic the natural LH peak and trigger ovulation. An injection of human chorionic gonadotrophin (hCG) is used to trigger ovulation at the end of stimulation with FSH or a mixture of FSH and LH. CG may also be used together with FSH during stimulation for OI and COH, in order to provide LH-activity during stimulation in patients in which LH-activity is desirable, such as those mentioned above.

FSH, LH and CG are members of the heterodimer, glycoprotein hormone family that also includes thyroid stimulating hormone (TSH). The members of this family are heterodimers, comprising an α - and a β -subunit. The subunits are held together by noncovalent interactions. The human FSH (hFSH) heterodimer consists of (i) a mature 92 amino acid glycoprotein alpha subunit, which also is common to the other human family members (i.e., chorionic gonadotrophin ("CG"), luteinising hormone ("LH") and thyroid stimulating hormone ("TSH")); and (ii) a mature 111 amino acid beta subunit that is unique to FSH². The human LH heterodimer consists of (i) the mature 92 amino acid glycoprotein alpha subunit; and (ii) a mature 112 beta subunit that is unique to LH³. The alpha and beta subunits of the glycoproteins may be prone to dissociate in formulations, due to interaction with a preservative, surfactant and other excipients. Dissociation of the subunits leads to loss of biological potency⁴.

FSH is formulated for intramuscular (IM) or subcutaneous (SC) injection. FSH is supplied in lyophilised (solid) form in vials or ampoules of 75 IU/vial and 150 IU/vial with a shelf life of one and a half to two years when stored at 2-25°C. A solution for injection is formed by reconstituting the lyophilised product with water for injection (WFI). For ovulation induction or controlled ovarian hyperstimulation, daily injections with starting doses of 75 IU to 600 IU are recommended for up to about ten days. Depending on the patient's response, up to three cycles of treatment with increasing doses of FSH can be used. With lyophilised formulations, the patient is required to reconstitute a new vial of lyophilised material with diluent and administer it immediately after reconstitution on a daily basis [Package insert N1700101A,

published in February 1996, for Fertinex™ (urofolitropin for injection, purified) for subcutaneous injection, by Serono Laboratories, Inc., Randolph, MA].

FSH has also been formulated in both single-dose and multi-dose liquid formats, in vials, or ampoules. Single dose formats must remain stable and potent in storage
5 prior to use. Multi-dose formats must not only remain stable and potent in storage prior to use, but must also remain stable, potent and relatively free of bacteria over the multiple-dose use regimen administration period, after the seal of the ampoule has been compromised. For this reason, multi-dose formats often contain a bacteriostatic agent.

10 LH is formulated for intramuscular (IM) or subcutaneous (SC) injection. LH is supplied in lyophilised (solid) form in vials or ampoules of 75 IU/vial with a shelf life of one and a half to two years when stored at 2-25°C. A solution for injection is formed by reconstituting the lyophilised product with water for injection (WFI). For ovulation
15 induction or controlled ovarian hyperstimulation, In conjunction with FSH, daily injections with starting doses of 75 IU to 600 IU LH are recommended for up to about ten days.

20 EP 0 618 808 (Applied Research Systems ARS Holding N.V.) discloses a pharmaceutical composition comprising a solid intimate mixture of gonadotrophin and a stabilising amount of sucrose alone or in combination with glycine.

EP 0 814 841 (Applied Research Systems ARS Holding N.V.) discloses a stable, liquid pharmaceutical composition comprising recombinant human chorionic
25 gonadotrophin (hCG) and a stabilizing amount of mannitol.

EP 0 448 146 (AKZO N.V.) discloses a stabilized gonadotrophin containing lyophilisate comprising one part by weight of a gonadotrophin; and 200 to 10,000 parts by weight of a dicarboxylic acid salt stabilizer associated with the
30 gonadotrophin.

EP 0 853 945 (Akzo Nobel N.V.) discloses a liquid gonadotrophin-containing formulation characterised in that the formulation comprises a gonadotrophin and stabilising amounts of a polycarboxylic acid or a salt thereof and of a thioether
35 compound.

WO 00/04913 (Eli Lilly and Co.) discloses a formulation comprising FSH or an FSH variant, containing an alpha and beta subunit, and a preservative selected from the group consisting of phenol, m-cresol, p-cresol, o-cresol, chlorocresol, benzyl alcohol, alkylparaben (methyl, ethyl, propyl, butyl and the like), benzalkonium chloride, benzethonium chloride, sodium dehydroacetate and thimerosal, or mixtures thereof in an aqueous diluent.

There remains a need for stable liquid formulations of FSH or FSH variants, and mixtures of FSH and LH, either for single dose or multiple dose administration.

Summary of the invention

It is an object of the invention to provide new freeze dried as well as liquid formulations of FSH or FSH variants, LH or LH variants, to provide methods for their preparation, and methods for their pharmaceutical or veterinary use in the treatment of fertility disorders.

It is a further object of the invention to provide new freeze dried and liquid formulations of mixtures of FSH and LH, to provide methods for their preparation, and methods for their pharmaceutical or veterinary use in the treatment of fertility disorders.

In a first aspect, the invention provides a freeze dried and liquid pharmaceutical composition comprising FSH or a variant thereof, and a surfactant selected from block copolymers of ethylene oxide and propylene oxide, preferably Pluronic® F77, Pluronic F87, Pluronic F88 and Pluronic F68.

In a second aspect, the invention provides a method for manufacturing a liquid pharmaceutical composition comprising forming a solution of FSH or a variant thereof, and a surfactant selected from block copolymers of ethylene oxide and propylene oxide, preferably Pluronic® F77, Pluronic F87, Pluronic F88 and Pluronic F68 and WFI.

In a third aspect, the invention provides a method for manufacturing a packaged pharmaceutical composition comprising dispensing a solution comprising FSH, and a surfactant selected from block copolymers of ethylene oxide and propylene oxide,

preferably Pluronic® F77, Pluronic F87, Pluronic F88 and Pluronic F68, into a container.

5 In a fourth aspect, the invention provides an article of manufacture for human pharmaceutical use, comprising a vial comprising a solution of FSH or an FSH variant, and a surfactant selected from block copolymers of ethylene oxide and propylene oxide, preferably Pluronic® F77, Pluronic F87, Pluronic F88 and Pluronic F68 and written material stating that such solution may be held over a period of at or about twenty-four hours or greater after the first use.

10 In a fifth aspect, the invention provides a freeze dried and liquid pharmaceutical composition comprising FSH and LH, and a surfactant selected from block copolymers of ethylene oxide and propylene oxide, preferably Pluronic® F77, Pluronic F87, Pluronic F88 and Pluronic F68.

15 In a sixth aspect, the invention provides a method for manufacturing a freeze dried and liquid pharmaceutical composition comprising forming a solution of FSH and LH and a surfactant selected from selected from block copolymers of ethylene oxide and propylene oxide, preferably Pluronic® F77, Pluronic F87, Pluronic F88 and Pluronic F68.

20 In an seventh aspect, the invention provides a method for manufacturing a packaged pharmaceutical composition comprising dispensing a solution comprising FSH and LH, and a surfactant selected from block copolymers of ethylene oxide and propylene oxide, preferably Pluronic® F77, Pluronic F87, Pluronic F88 and Pluronic F68, into a container.

25 In an eighth aspect, the invention provides an article of manufacture for human pharmaceutical use, comprising a vial comprising a solution of FSH and LH, and a surfactant selected from block copolymers of ethylene oxide and propylene oxide, preferably Pluronic® F77, Pluronic F87, Pluronic F88 and Pluronic F68 and written material stating that such solution may be held over a period of at or about twenty-four hours or greater after the first use.

30 In a ninth aspect, the invention provides an article of manufacture for human pharmaceutical use, comprising a first container comprising freeze dried FSH or an

FSH variant, and a surfactant selected from block copolymers of ethylene oxide and propylene oxide, preferably Pluronic® F77, Pluronic F87, Pluronic F88 and Pluronic F68 and a second container comprising a solvent for reconstitution, preferably an aqueous solution containing a bacteriostatic, preferably m-cresol.

5

In an tenth aspect, the invention provides an article of manufacture for human pharmaceutical use, comprising a first container comprising freeze dried LH or an LH variant, and a surfactant selected from block copolymers of ethylene oxide and propylene oxide, preferably Pluronic® F77, Pluronic F87, Pluronic F88 and Pluronic F68 and a second container comprising a solvent for reconstitution, preferably an aqueous solution containing a bacteriostatic, preferably m-cresol.

10

In an eleventh aspect, the invention provides an article of manufacture for human pharmaceutical use, comprising a first container comprising freeze dried FSH as well as LH or an FSH or LH variant variant, and a surfactant selected from block copolymers of ethylene oxide and propylene oxide, preferably Pluronic® F77, Pluronic F87, Pluronic F88 and Pluronic F68 and a second container comprising a solvent for reconstitution, preferably an aqueous solution with m-cresol.

15

20 Detailed description of the invention

Brief description of the drawings

Figure 1 shows the percentage of oxidised α -subunit in formulations of FSH containing Pluronic F68, methionine at 10 $\mu\text{g/ml}$ ("Meth 10 mcg/ml") and 100 $\mu\text{g/ml}$ ("Meth 100 mcg/ml") versus a formulation with no methionine ("No methionine"), at

25

time 0, 1 week and 2 weeks.

The liquid and freeze dried FSH or FSH and LH formulations of the invention have improved or more suitable properties or stability, and are useful for infertility treatment in women and/or men. These formulations and articles of manufacture are additionally suitable for use in injectable and alternative delivery systems, e.g., but not limited to, nasal, pulmonary, transmucosal, transdermal, oral, subcutaneous, intramuscular or parenteral sustained release. In a particularly preferred embodiment the formulations of the invention are for subcutaneous and/or intramuscular injection. The FSH or FSH and LH variant solutions and formulations provided may also have increased *in vivo* potency over time compared to known commercial products, by preventing or reducing loss of activity or stability, or by

30

35

improving any aspect of the effectiveness or desirability of administration, e.g., by at least one of mode, frequency, dosage, comfort, ease of use, biological activity *in vitro* or *in vivo*, and the like.

- 5 Follicle stimulating hormone, or FSH, as used herein refers to the FSH produced as a full-length mature protein which includes, but is not limited to human FSH or "hFSH", whether produced recombinantly or isolated from human sources, such as the urine of postmenopausal women. The protein sequence of the human glycoprotein alpha subunit is provided in SEQ ID NO: 1, and the protein sequence of the human FSH
- 10 beta subunit is given in SEQ ID NO:2.

- The expression "FSH variant" is meant to encompass those molecules differing in amino acid sequence, glycosylation pattern or in inter-subunit linkage from human FSH but exhibiting FSH-activity. Examples include CTP-FSH, a long-acting modified
- 15 recombinant FSH, consisting of the wild type α -subunit and a hybrid β -subunit in which the carboxy terminal peptide of hCG has been fused to the C-terminal of the β -subunit of FSH, as described in LaPolt *et al.*; Endocrinology; 1992, 131, 2514-2520; or Klein *et al.*; Development and characterization of a long-acting recombinant hFSH agonist; Human Reprod. 2003, 18, 50-56]. Also included is single chain CTP-
- 20 FSH, a single chain molecule, consisting of the following sequences (from N-terminal to C-terminal):

β FSH	β hCG-CTP(113-145)	α FSH
-------------	--------------------------	--------------

- 25 wherein β FSH signifies the β -subunit of FSH, β hCG CTP (113-145) signifies the carboxy terminal peptide of hCG and α FSH signifies the α -subunit of FSH, as described by Klein *et al.*⁵ Other examples of FSH variants include FSH molecules having additional glycosylation sites incorporated in the α - and/or β -subunit, as
- 30 disclosed in WO 01/58493 (Maxygen), particularly as disclosed in claims 10 and 11 of WO 01/58493, and FSH molecules with intersubunit S-S bonds, as disclosed in WO 98/58957.

- The FSH variants referred to herein also include the carboxy terminal deletions of the beta subunit that are shorter than the full length mature protein of SEQ ID NO:2.
- 35 Carboxy terminal deletions of the human beta subunit are provided in SEQ IDS NOS:

3, 4, and 5. It is understood that the carboxy terminal variants of the beta chain form dimers with a known alpha subunit to form an FSH variant heterodimer.

5 FSH heterodimers or FSH variant heterodimers can be produced by any suitable method, such as recombinantly, by isolation or purification from natural sources as may be the case, or by chemical synthesis, or any combination thereof.

10 The use of the term "recombinant" refers to preparations of FSH, LH or FSH and LH variants that are produced through the use of recombinant DNA technology (see for example WO 85/01958). The sequences for genomic and cDNA clones of FSH are known for the alpha and beta subunits of several species⁶. One example of a method of expressing FSH or LH using recombinant technology is by transfection of eukaryotic cells with the DNA sequences encoding an alpha and beta subunit of FSH or LH, whether provided on one vector or on two vectors with each subunit having a
15 separate promoter, as described in European patent nos. EP 0 211 894 and EP 0 487 512. Another example of the use of recombinant technology to produce FSH or LH is by the use of homologous recombination to insert a heterologous regulatory segment in operative connection to endogenous sequences encoding the subunits of FSH or LH, as described in European patent no. EP 0 505 500 (Applied Research
20 Systems ARS Holding NV).

The FSH or FSH variant used in accordance with the present invention may be produced not only by recombinant means, including from mammalian cells, but also may be purified from other biological sources, such as from urinary sources.
25 Acceptable methodologies include those described in Hakola, K. Molecular and Cellular Endocrinology, 127:59-69, 1997; Keene, et al., J. Biol. Chem., 264:4769-4775, 1989; Cerpa-Poljak, et al., Endocrinology, 132:351-356, 1993; Dias, et al., J. Biol. Chem., 269:25289-25294, 1994; Flack, et al., J. Biol. Chem., 269:14015-14020, 1994; and Valove, et al., Endocrinology, 135:2657-2661, 1994, U.S. Patent
30 3,119,740 and US Patent no. 5,767,067.

Luteinising hormone, or LH, as used herein refers to the LH produced as a full-length mature protein, which includes, but is not limited to human LH or "hLH", whether
35 produced recombinantly or isolated from human sources, such as the urine of postmenopausal women. The protein sequence of the human glycoprotein alpha

subunit is provided in SEQ ID NO: 1, and the protein sequence of the human LH beta subunit⁷ is given in SEQ ID NO: 6. In a preferred embodiment the LH is recombinant.

5 The expression "LH variant" is meant to encompass those molecules differing in amino acid sequence, glycosylation pattern or in inter-subunit linkage from human LH but exhibiting LH-activity.

10 LH heterodimers or LH variant heterodimers can be produced by any suitable method, such as recombinantly, by isolation or purification from natural sources as may be the case, or by chemical synthesis, or any combination thereof.

15 The term "administer" or "administering" means to introduce a formulation of the present invention into the body of a patient in need thereof to treat a disease or condition.

The term "patient" means a mammal that is treated for a disease or condition. Patients are of, but not limited to, the following origin, human, ovine, porcine, equine, bovine, rabbit and the like.

20 The term "potency" in relation to FSH activity, refers to the ability of an FSH formulation or a mixed formulation, to elicit biological responses associated with FSH, such as ovarian weight gain in the Steelman-Pohley assay⁸, or follicular growth in a female patient. Follicular growth in a female patient can be evaluated by ultrasound, for example, in terms of the number of follicles having a mean diameter
25 of at or about 16 mm on day 8 of stimulation. Biological activity is evaluated with respect to an accepted standard for FSH.

The term "potency" in relation to LH activity, refers to the ability of an LH formulation or a mixed formulation, to elicit biological responses associated with LH, such as
30 seminal vesicle weight gain method.⁹ Biological activity of LH is evaluated with respect to an accepted standard for LH.

35 The term "aqueous diluent" refers to a liquid solvent that contains water. Aqueous solvent systems may be consist solely of water, or may consist of water plus one or more miscible solvents, and may contain dissolved solutes such as sugars, buffers, salts or other excipients. The more commonly used non-aqueous solvents are the

short-chain organic alcohols, such as, methanol, ethanol, propanol, short-chain ketones, such as acetone, and poly alcohols, such as glycerol.

5 An "isotonicity agent" is a compound that is physiologically tolerated and imparts a suitable tonicity to a formulation to prevent the net flow of water across cell membranes that are in contact with the formulation. Compounds such as glycerin, are commonly used for such purposes at known concentrations. Other suitable isotonicity agents include, but are not limited to, amino acids or proteins (e.g., glycine or albumin), salts (e.g., sodium chloride), and sugars (e.g., dextrose, sucrose and lactose).

15 The term "bacteriostatic" or "bacteriostatic agent" refers to a compound or compositions added to a formulation to act as an anti-bacterial agent. A preserved FSH or FSH variant or FSH and LH containing formulation of the present invention preferably meets statutory or regulatory guidelines for preservative effectiveness to be a commercially viable multi-use product, preferably in humans. Examples of bacteriostatics include phenol, *m*-cresol, *p*-cresol, *o*-cresol, chlorocresol, benzyl alcohol, alkylparaben (methyl, ethyl, propyl, butyl and the like), benzalkonium chloride, benzethonium chloride, sodium dehydroacetate and thimerosal.

20 The term "buffer" or "physiologically-acceptable buffer" refers to solutions of compounds that are known to be safe for pharmaceutical or veterinary use in formulations and that have the effect of maintaining or controlling the pH of the formulation in the pH range desired for the formulation. Acceptable buffers for controlling pH at a moderately acidic pH to a moderately basic pH include, but are not limited to, such compounds as phosphate, acetate, citrate, arginine, TRIS, and histidine. "TRIS" refers to 2-amino-2-hydroxymethyl-1,3,-propanediol, and to any pharmacologically acceptable salt thereof. Preferable buffers are phosphate buffers with saline or an acceptable salt.

25 30 The term "phosphate buffer" refers to solutions containing phosphoric acid or salts thereof, adjusted to a desired pH. Generally phosphate buffers are prepared from phosphoric acid, or a salt of phosphoric acid, including but not limited to sodium and potassium salts. Several salts of phosphoric acid are known in the art, such as sodium and potassium monobasic, dibasic, and tribasic salts of the acid. Salts of phosphoric acid are also known to occur as hydrates of the occurring salt. Phosphate

buffers may cover a range of pHs, such as from about pH 4 to about pH 10, and preferred ranges from about pH 5 to about pH 9, and a most preferred range of at or about 6.0 to at or about 8.0, most preferably at or about pH 7.0.

- 5 The term "vial" refers broadly to a reservoir suitable for retaining FSH in solid or liquid form in a contained sterile state. Examples of a vial as used herein include ampoules, cartridges, blister packages, or other such reservoir suitable for delivery of the FSH to the patient via syringe, pump (including osmotic), catheter, transdermal patch, pulmonary or transmucosal spray. Vials suitable for packaging products for
10 parenteral, pulmonary, transmucosal, or transdermal administration are well known and recognized in the art.

- The term "stability" refers to the physical, chemical, and conformational stability of FSH and LH in the formulations of the present invention (including maintenance of
15 biological potency). Instability of a protein formulation may be caused by chemical degradation or aggregation of the protein molecules to form higher order polymers, by dissociation of the heterodimers into monomers, deglycosylation, modification of glycosylation, oxidation (particularly of the α -subunit) or any other structural modification that reduces at least one biological activity of an FSH polypeptide
20 included in the present invention.

- A "stable" solution or formulation, is one wherein the degree of degradation, modification, aggregation, loss of biological activity and the like, of proteins therein is acceptably controlled, and does not increase unacceptably with time. Preferably the
25 formulation retains at least at or about 80% of the labelled FSH activity and at least at or about 80% of the labelled LH activity over a period of 6 months at a temperature of at or about 2-8°C, more preferably at or about 2-8°C, more preferably at or about 4-5°C. FSH activity can be measured using the Steelman-Pohley ovarian weight gain bioassay⁵. LH activity can be measured using the seminal vesicle weight gain
30 bioassay¹⁰.

- The term "treating" refers to the administration, follow up, management and/or care of a patient for which FSH and/or LH administration is desirable for the purpose of follicle or testicular stimulation or any other physiological response regulated by FSH
35 and/or LH. Treating can thus include, but is not limited to, the administration of FSH and/or LH for the induction or improvement of sperm quality, stimulation of

testosterone release in the male, or follicular development or for ovulation induction in the female.


5 The expression "multi-dose use" is intended to include the use of a single vial, ampoule or cartridge of an FSH formulation or a formulation of FSH and LH for more than one injection, for example 2, 3, 4, 5, 6 or more injections. The injections are preferably made over a period of at least at or about 12 hours, 24 hours, 48 hours, etc., preferably up to a period of at or about 12 days. The injections may be spaced in time, for example, by a period of 6, 12, 24, 48 or 72 hours.

10 A "salt" of a protein is an acid or base addition salt. Such salts are preferably formed between any one or more of the charged groups in the protein and any one or more physiologically acceptable, non-toxic cations or anions. Organic and inorganic salts include, for example, those prepared from acids such as hydrochloric, sulphuric, sulfonic, tartaric, fumaric, hydrobromic, glycolic, citric, maleic, phosphoric, succinic, 15 acetic, nitric, benzoic, ascorbic, p-toluenesulfonic, benzenesulfonic, naphthalenesulfonic, propionic, carbonic, and the like, or for example, ammonium, sodium, potassium, calcium, or magnesium.

20 The inventors have found that by formulating FSH and mixtures of FSH and LH with a surfactant selected from block copolymers of ethylene oxide and propylene oxide, preferably Pluronic® F77, Pluronic F87, Pluronic F88 and Pluronic® F68, particularly preferably Pluronic F68 (BASF, Pluronic F68 is also known as Poloxamer 188) they obtain stable formulations that minimise the loss of active principle (FSH or FSH and 25 LH) caused by adsorption on the surfaces of the vial and/or delivery device (e.g. syringe, pump, catheter, etc.).

The inventors have further found that by formulating FSH and mixtures of FSH and LH with a surfactant selected from block copolymers of ethylene oxide and propylene 30 oxide, preferably Pluronic® F77, Pluronic F87, Pluronic F88 and Pluronic® F68, particularly preferably Pluronic F68 (BASF, Pluronic F68 is also known as Poloxamer 188) they obtain a stable formulation that avoids the problem of precipitation in the presence of a bacteriostatic agent, such as *m*-cresol and phenol. Precipitation, resulting in the formation of turbid or milky solutions occurs when TWEEN 20 is used 35 with *m*-cresol or phenol.

The Pluronic surfactants are block copolymers of ethylene oxide (EO) and propylene oxide (PO). The propylene oxide block (PO) is sandwiched between two ethylene oxide (EO) blocks.



5

Pluronic surfactants are synthesised in a two-step process:

1. A hydrophobe of the desired molecular weight is created by the controlled addition of propylene oxide to the two hydroxyl groups of propylene glycol; and
2. Ethylene oxide is added to sandwich the hydrophobe between hydrophilic groups.

In Pluronic® F77, the percentage of polyoxyethylene (hydrophile) is 70%, and the molecular weight of the hydrophobe (polyoxypropylene) is approximately 2,306 Da.

- 15 In Pluronic F87, the percentage of polyoxyethylene (hydrophile) is 70%, and the molecular weight of the hydrophobe (polyoxypropylene) is approximately 2,644 Da.

In Pluronic F88, the percentage of polyoxyethylene (hydrophile) is 80%, and the molecular weight of the hydrophobe (polyoxypropylene) is approximately 2,644 Da.

- 20 In Pluronic F68, the percentage of polyoxyethylene (hydrophile) is 80%, and the molecular weight of the hydrophobe (polyoxypropylene) is approximately 1,967 Da.

Typical properties of Pluronic F77 are listed below:

- 25 Average Molecular Weight: 6600;
 Melt/pour point: 48°C;
 Physical Form @ 20°C : solid;
 Viscosity (Brookfield) cps: 480 [liquids at 25°C, pastes at 60°C and solids at 77°C];
 Surface tension, dynes/cm @ 25°C;
 0.1% Conc. : 47.0
 0.01% Conc. : 49.3
 0.001% Conc.: 52.8
 30 Interfacial tension, dynes/cm @ 25°C vs. Nujol;
 0.1% Conc. : 17.7
 0.01% Conc. : 20.8
 35 0.001% Conc. : 25.5

- Draves Wetting, Seconds 25°C
 1.0% Conc.: > 360
 0.1% Conc.: > 360
- Foam Height
- 5 Ross Miles, 0.1%, mm @ 50°C: 100
 Ross Miles, 0.1%, mm @ 26°C: 47
 Dynamic, 0.1%, mm @ 400 ml/min: > 600
- Cloud point in aqueous solution, °C
 1% Conc.: >100
 10% Conc.: >100
- 10 HLB (hydrophile-lipophile balance): 25
- Typical properties of Pluronic F87 are listed below:
 Average Molecular Weight: 7700;
 Melt/pour point: 49°C;
 Physical Form @ 20°C: solid;
 Viscosity (Brookfield) cps: 700 [liquids at 25°C, pastes at 60°C and solids at 77°C];
 Surface tension, dynes/cm @ 25°C;
 0.1% Conc.: 44.0
 0.01% Conc.: 47.0
 0.001% Conc.: 50.2
- 20 Interfacial tension, dynes/cm @ 25°C vs Nujol;
 0.1% Conc.: 17.4
 0.01% Conc.: 20.3
 0.001% Conc.: 23.3
- 25 Draves Wetting, Seconds 25°C
 1.0% Conc.: > 360
 0.1% Conc.: > 360
- Foam Height
- 30 Ross Miles, 0.1%, mm @ 50°C: 80
 Ross Miles, 0.1%, mm @ 26°C: 37
 Dynamic, 0.1%, mm @ 400 ml/min: > 600
- Cloud point in aqueous solution, °C
 1% Conc.: >100
 10% Conc.: >100
- 35 HLB (hydrophile-lipophile balance): 24
- Typical properties of Pluronic F88 are listed below:
 Average Molecular Weight: 11400;
 Melt/pour point: 54°C;
 Physical Form @ 20°C: solid;
 Viscosity (Brookfield) cps: 2300 [liquids at 25°C, pastes at 60°C and solids at 77°C];
 Surface tension, dynes/cm @ 25°C;
 0.1% Conc.: 48.5
 0.01% Conc.: 52.6
 0.001% Conc.: 55.7
- 45 Interfacial tension, dynes/cm @ 25°C vs Nujol;
 0.1% Conc.: 20.5
 0.01% Conc.: 23.3
 0.001% Conc.: 27.0
- 50 Draves Wetting, Seconds 25°C
 1.0% Conc.: > 360
 0.1% Conc.: > 360
- Foam Height

- Ross Miles, 0.1%, mm @ 50°C: 80
 Ross Miles, 0.1%, mm @ 26°C: 37
 Dynamic, 0.1%, mm @ 400 ml/min: > 600
 Cloud point in aqueous solution, °C
 1% Conc.: >100
 10% Conc.: >100
 HLB (hydrophile-lipophile balance): 28
- Typical properties of Pluronic F68 are listed below:
 Average Molecular Weight: 8400;
 Melt/pour point: 52°C;
 Physical Form @ 20°C : solid;
 Viscosity (Brookfield) cps: 1000 (liquids at 25°C, pastes at 60°C and solids at 77°C);
 Surface tension, dynes/cm @ 25°C;
 0.1% Conc. : 50.3
 0.01% Conc. : 51.2
 0.001% Conc.: 53.6
 Interfacial tension, dynes/cm @ 25°C vs Nujol;
 0.1% Conc. : 19.8
 0.01% Conc. : 24.0
 0.01% Conc. : 26.0
 Draves Wetting, Seconds 25°C
 1.0% Conc.: > 360
 0.1% Conc.: > 360
 Foam Height
 Ross Miles, 0.1%, mm @ 50°C: 35
 Ross Miles, 0.1%, mm @ 26°C: 40
 Dynamic, 0.1%, mm @ 400 ml/min: > 600
 Cloud point in aqueous solution, °C
 1% Conc.: >100
 10% Conc.: >100
 HLB (hydrophilelipophile balance): 29
- Other polymers having properties similar to those listed above may also be used in the formulations of the invention. The preferred surfactant is Pluronic F68, and surfactants having similar properties.
- Pluronic, particularly Pluronic F68, is preferably present in the formulation at a concentration that is sufficient to maintain FSH and/or LH stability over the desired storage period (for example 6 to 12 to 24 months), and also at a concentration that is sufficient to prevent protein losses due to adsorption on surfaces, such as the vial, ampoule or cartridge or the syringe.
- Preferably the concentration of Pluronic, particularly Pluronic F68, in liquid formulations is at or about 0.01 mg/ml to at or about 1 mg/ml, more preferably at or about 0.05 mg/ml to at or about 0.5 mg/ml, more particularly preferably at or about 0.2 mg/ml to at or about 0.4 mg/ml, most preferably at or about 0.1 mg/ml.

The follicle-stimulating hormone (FSH) within the freeze-dried formulation is preferably present at a concentration (w/w) of at or about 0.1 to 10 µg/mg of the total formulation. In one embodiment, the follicle-stimulating hormone (FSH) is present at
5 a concentration of at or about 0.3 to 5 µg/mg of the total formulation. In a further embodiment the follicle-stimulating hormone (FSH) is present at a concentration of at or about 0.37 to 2 µg/mg of the total formulation.

The luteinising hormone (LH) within the freeze-dried formulation is preferably present
10 at a concentration of at or about 0.1 to 3 µg/mg of the total formulation. In one embodiment, the luteinising hormone (LH) is present at a concentration of at or about 0.1 to 1 µg/mg of the total formulation. In a further embodiment, the luteinising hormone (LH) is present at a concentration of at or about 0.1 to 0.6 µg/mg of the total formulation.

15 In the liquid formulations – including the reconstituted formulations – comprising FSH, preferably the concentration of FSH in the formulation is at or about 150 IU/ml to at or about 2,000 IU/ml, more preferably at or about 300 IU/ml to at or about 1,500 IU/ml, more particularly preferably at or about 450 to at or about 750, most preferably at or
20 about 600 IU/ml.

In the liquid formulations – including the reconstituted formulations – comprising LH, preferably the LH concentration in the formulation is at or about 50 IU/ml to at or about 2,000 IU/ml, more preferably at or about 150 to at or about 1,500 IU/ml, more
25 particularly preferably at or about 300 IU/ml to at or about 750 IU/ml, particularly preferably 625 IU/ml.

In formulations comprising both FSH and LH, the ratio of FSH to LH (FSH:LH, IU:IU, FSH measured with rat ovarian weight gain assay and LH measured with rat seminal vesicle weight gain assay) is preferably within the range of at or about 6:1 to at or
30 about 1:6, more preferably at or about 4:1 to at or about 1:2, more particularly preferably at or about 3:1 to at or about 1:1. Particularly preferred ratios are 1:1 and 2:1.

In the freeze dried formulations, the surfactant, e.g. Pluronic F 68, is preferably present at a concentration of at or about 0.001 to at or about 0.1 mg per mg of the
35 total formulation, more preferably at or about 0.01 to at or about 0.075 mg/mg.

Preferably the concentration of Pluronic, particularly Pluronic F68, in the reconstituted formulations is at or about 0.01 mg/ml to at or about 1 mg/ml, more preferably at or about 0.05 mg/ml to at or about 0.5 mg/ml, more particularly preferably at or about 0.2 mg/ml to at or about 0.4 mg/ml, most preferably at or about 0.1 mg/ml.

Preferably the FSH and LH are produced recombinantly, particularly preferably they are produced in Chinese hamster ovary cells transfected with a vector or vectors comprising DNA coding for the human glycoprotein alpha-subunit and the beta-subunit of FSH or LH. DNA encoding the alpha and beta-subunits may be present on the same or different vectors.

Recombinant FSH and LH have several advantages over their urinary counterparts. Culture and isolation techniques using recombinant cells permit consistency between batches. In contrast, urinary FSH and LH vary greatly from batch to batch in such characteristics as purity, glycosylation pattern, sialylation and oxidation of the subunits. Due to greater batch-to-batch consistency and purity of recombinant FSH and LH, the hormones can be readily identified and quantified using techniques such as isoelectric focussing (IEF). The ease with which recombinant FSH and LH can be identified and quantified permits the filling of vials by mass of hormone (fill-by-mass) rather than filling by bioassay.

Preferably formulations of FSH of the present invention have pH between at or about 6.0 and at or about 8.0, more preferably at or about 6.8 to at or about 7.8, including about pH 7.0, pH 7.2, and 7.4. A preferred buffer is phosphate, with preferred counterions being sodium or potassium ions. Phosphate saline buffers are well known in the art, such as Dulbecco's Phosphate buffered saline. Buffer concentrations in total solution can vary between at or about 5mM, 9.5mM, 10mM, 50mM, 100mM, 150mM, 200mM, 250mM, and 500mM. Preferably the buffer concentration is at or about 10mM. Particularly preferred is a buffer 10 mM in phosphate ions with a pH of 7.0.

Preferably formulations of mixtures of FSH and LH of the present invention have pH between at or about 6.0 and at or about 9.0, more preferably at or about 6.8 to at or about 8.5, including about pH 7.0, pH 8.0, and 8.2, most preferably at or about pH 8.0.

The invention is directed to liquid formulations as well as freeze dried (lyophilised) formulations that may be reconstituted, in which the solvent (also for reconstitution) is water for injection. Liquid formulations may be single dose or multi-dose. Those liquid as well as freeze dried FSH and/or LH formulations of the invention that are intended for multi-dose use preferably comprise a bacteriostatic, such as phenol, *m*-cresol, *p*-cresol, *o*-cresol, chlorocresol, benzyl alcohol, alkylparaben (methyl, ethyl, propyl, butyl and the like), thymol, benzalkonium chloride, benzethonium chloride, sodium dehydroacetate and thimerosal. Particularly preferred are phenol, benzyl alcohol and *m*-cresol, more preferred are phenol and *m*-cresol, most preferred is *m*-cresol. The bacteriostatic agent is used in an amount that will yield a concentration that is effective to maintain the formulation essentially bacteria free (suitable for injection) over the multi-dose injection period, which may be at or about 12 or 24 hours to at or about 12 or 14 days, preferably at or about 6 to at or about 12 days. The bacteriostatic is preferably present in a concentration of at or about 0.1% (mass bacteriostatic/mass of solvent) to at or about 2.0%, more preferably at or about 0.2% to at or about 1.0%. In the case of benzyl alcohol, particularly preferred is a concentration of 0.9%. In the case of phenol, particularly preferred is at or about 0.5%. In the case of *m*-cresol, particularly preferred is a concentration of at or about 0.3 % (e.g. at or about 3 mg/ml in WFI).

In a preferred embodiment, the invention provides a liquid pharmaceutical composition, preferably for multi-dose use, comprising FSH or a variant thereof, a surfactant selected from Pluronic® F77, Pluronic F87, Pluronic F88 and Pluronic F68, and a bacteriostatic selected from *m*-cresol and phenol, preferably *m*-cresol.

In a further preferred embodiment, the invention provides a liquid pharmaceutical composition, preferably for multi-dose use, comprising LH, a surfactant selected from Pluronic® F77, Pluronic F87, Pluronic F88 and Pluronic F68, and a bacteriostatic selected from *m*-cresol and phenol, preferably *m*-cresol.

In a further preferred embodiment, the invention provides a liquid pharmaceutical composition, preferably for multi-dose use, comprising FSH and LH, a surfactant selected from Pluronic® F77, Pluronic F87, Pluronic F88 and Pluronic F68, and a bacteriostatic selected from *m*-cresol and phenol, preferably *m*-cresol. Preferably the FSH and LH are present in a ratio (FSH:LH) of at or about 2:1 to at or about 1:1.

In a further preferred embodiment, the invention provides a method for manufacturing a liquid pharmaceutical composition, preferably for multi-dose use, comprising forming an aqueous solution of FSH or a variant thereof, a surfactant selected from Pluronic® F77, Pluronic F87, Pluronic F88 and Pluronic F68, and a bacteriostatic selected from m-cresol and phenol, preferably m-cresol, and WFI.

In a further preferred embodiment, the invention provides a method for manufacturing a liquid pharmaceutical composition, preferably for multi-dose use, comprising forming an aqueous solution of LH, a surfactant selected from Pluronic® F77, Pluronic F87, Pluronic F88 and Pluronic F68, and a bacteriostatic selected from m-cresol and phenol, preferably m-cresol, and WFI.

In a further preferred embodiment, the invention provides a method for manufacturing a liquid pharmaceutical composition, preferably for multi-dose use, comprising forming an aqueous solution of FSH and LH, a surfactant selected from Pluronic® F77, Pluronic F87, Pluronic F88 and Pluronic F68, and a bacteriostatic selected from m-cresol and phenol, preferably m-cresol, and WFI.

In yet another preferred embodiment, the invention provides a method for manufacturing a packaged pharmaceutical composition comprising dispensing a solution comprising FSH, a surfactant selected from Pluronic® F77, Pluronic F87, Pluronic F88 and Pluronic F68, and a bacteriostatic selected from m-cresol and phenol, preferably m-cresol.

In yet another preferred embodiment, the invention provides a method for manufacturing a packaged pharmaceutical composition comprising dispensing a solution comprising FSH and LH, a surfactant selected from Pluronic® F77, Pluronic F87, Pluronic F88 and Pluronic F68, and a bacteriostatic selected from m-cresol and phenol, preferably m-cresol.

In yet another preferred embodiment, the invention provides an article of manufacture for human pharmaceutical use, comprising a vial comprising a solution of FSH or an FSH variant, a surfactant selected from Pluronic® F77, Pluronic F87, Pluronic F88 and Pluronic F68, and a bacteriostatic selected from m-cresol and phenol, preferably m-cresol, and written material stating that such solution may be held over a period of at or about twenty-four hours or greater after the first use.

Preferably the written material states that the solution may be held up to at or about 12 or 14 days after the first use.

In yet another preferred embodiment, the invention provides an article of manufacture for human pharmaceutical use, comprising a vial comprising a solution of FSH and LH, a surfactant selected from Pluronic® F77, Pluronic F87, Pluronic F88 and Pluronic F68, and a bacteriostatic selected from *m*-cresol and phenol, preferably *m*-cresol, and written material stating that such solution may be held over a period of at or about twenty-four hours or greater after the first use. Preferably the written material states that the solution may be held up to at or about 12 or 14 days after the first use.

In a particularly preferred embodiment, the formulation comprises *m*-cresol and Pluronic F68. The inventors have surprisingly found that formulations comprising Pluronic F68 do not precipitate in the presence of *m*-cresol, a problem observed with other surfactants.

Before the first use, that is before the seal of the vial ampoule or cartridge has been broken, the formulations of the invention may be kept for at least at or about 6 months, 12 months or 24 months. Under preferred storage conditions, before the first use, the formulations are kept away from bright light (preferably in the dark), at temperatures of at or about 2-8°C, more preferably at or about 4-5°C.

In a specific embodiment, the invention provides a freeze dried formulation for reconstitution, preferably for multi-dose use, comprising FSH or a variant thereof and a surfactant selected from Pluronic® F77, Pluronic F87, Pluronic F88 and Pluronic F68, preferably Pluronic F68.

In a further specific embodiment, the invention provides a freeze dried formulation for reconstitution, preferably for multi-dose use, comprising LH, a surfactant selected from Pluronic® F77, Pluronic F87, Pluronic F88 and Pluronic F68, preferably Pluronic F68.

In a further specific embodiment, the invention provides a freeze dried formulation, preferably for multi-dose use, comprising FSH and LH, a surfactant selected from Pluronic® F77, Pluronic F87, Pluronic F88 and Pluronic F68, preferably Pluronic F68.

Preferably the FSH and LH are present in a ratio (FSH:LH) of at or about 2:1 to at or about 1:1.

5 In a further specific embodiment, the invention provides a method for manufacturing a freeze dried formulation, preferably for multi-dose use after reconstitution, comprising forming a mixture of FSH or a variant thereof with a surfactant selected from Pluronic® F77, Pluronic F87, Pluronic F88 and Pluronic F68, and subjecting said mixture to lyophilisation.

10 In a further specific embodiment, the invention provides a method for manufacturing a freeze dried formulation, preferably for multi-dose use after reconstitution, comprising forming a mixture of LH with a surfactant selected from Pluronic® F77, Pluronic F87, Pluronic F88 and Pluronic F68, and subjecting said mixture to lyophilisation.

15 In a further specific embodiment, the invention provides a method for manufacturing a freeze dried formulation, preferably for multi-dose use after reconstitution, comprising forming a mixture of FSH and LH as well as a surfactant selected from Pluronic® F77, Pluronic F87, Pluronic F88 and Pluronic F68, and subjecting said mixture to lyophilisation.

20 In yet another preferred embodiment, the invention provides a method for manufacturing a packaged pharmaceutical composition comprising dispensing a freeze dried mixture comprising FSH and a surfactant selected from Pluronic® F77, Pluronic F87, Pluronic F88 and Pluronic F68.

25 In yet another preferred embodiment, the invention provides a method for manufacturing a packaged pharmaceutical composition comprising dispensing a freeze dried mixture comprising LH and a surfactant selected from Pluronic® F77, Pluronic F87, Pluronic F88 and Pluronic F68 into a container.

30 In yet another preferred embodiment, the invention provides a method for manufacturing a packaged pharmaceutical composition comprising dispensing a freeze dried mixture comprising FSH as well as LH and a surfactant selected from Pluronic® F77, Pluronic F87, Pluronic F88 and Pluronic F68 into a container.

In yet another preferred embodiment, the invention provides an article of manufacture for human pharmaceutical use, comprising a first container or vial comprising freeze dried FSH or an FSH variant and a surfactant selected from Pluronic® F77, Pluronic F87, Pluronic F88 and Pluronic F68. A second container or vial contains a diluent for reconstitution, preferably water and a bacteriostatic selected from m-cresol and phenol, preferably m-cresol.

In yet another preferred embodiment, the invention provides an article of manufacture for human pharmaceutical use, comprising a first container or vial comprising freeze dried LH or an LH variant and a surfactant selected from Pluronic® F77, Pluronic F87, Pluronic F88 and Pluronic F68. A second container or vial contains a diluent for reconstitution, preferably water and a bacteriostatic selected from m-cresol and phenol, preferably m-cresol.

In yet another preferred embodiment, the invention provides an article of manufacture for human pharmaceutical use, comprising a first container or vial comprising freeze dried FSH or an FSH variant as well as LH or an LH variant and a surfactant selected from Pluronic® F77, Pluronic F87, Pluronic F88 and Pluronic F68. A second container or vial contains a diluent for reconstitution, preferably water and a bacteriostatic selected from m-cresol and phenol, preferably m-cresol.

In a particularly preferred embodiment, the solvent for reconstitution comprises m-cresol. The inventors have found that freeze dried formulations comprising Pluronic F68 do not precipitate when reconstituted with a diluent containing m-cresol, a problem observed with other surfactants, e.g. Tween.

The freeze dried formulations of the invention may be kept for at least at or about 6 months, 12 months or 24 months. Under preferred storage conditions, before the first use, the formulations are kept away from bright light (preferably in the dark), at temperatures of at or about 25, preferably of at or about 2-8°C, more preferably at or about 4-5°C.

After the first use of a liquid or a reconstituted multi-dose formulation it may be kept and used for at least at or about 24 hours, preferably at least at or about 4, 5 or 6 days, more preferably for up to 12 or 14 days. After the first use the formulation is preferably stored at below room temperature (i.e. below at or about 25°C), more

preferably below at or about 10°C, more preferably at or about 2-8°C, most preferably at or about 5-0°C.

5 Preferably the formulations of the invention contain an antioxidant, such as methionine, sodium bisulfite, salts of ethylenediaminetetraacetic acid (EDTA), butylated hydroxytoluene (BHT), and butylated hydroxy anisole (BHA). Most preferred is methionine. The antioxidant prevents oxidation of FSH and LH (particularly the α -subunit).

10 Methionine in the liquid and/or reconstituted formulation is preferably present at a concentration of at or about 0.01 to at or about 1.0 mg/ml, more preferably at or about 0.05 to at or about 0.5 mg/ml, most preferably at or about 0.1 mg/ml.

15 Preferably the formulations of the invention contain a mono- or disaccharide or a sugar alcohol as stabiliser and tonicity adjusting agent, such as sucrose, dextrose, lactose, mannitol and/or glycerol. Most preferred is sucrose, preferably at a concentration of at or about 60 mg/ml.

20 As noted above, the invention provides liquid formulations for single use and multi-dose use, containing a bacteriostatic, or to which a bacteriostatic is added when the formulation is reconstituted. The formulations of the invention are suitable for pharmaceutical or veterinary use.

25 As noted above, in a preferred embodiment, the invention provides an article of manufacture, comprising packaging material and a vial comprising a solution of FSH or an FSH variant, LH, or FSH and LH, Pluronic F68 and a bacteriostatic selected from phenol and *m*-cresol, optionally with buffers and/or other excipients, in an aqueous diluent, wherein said packaging material comprises written material which indicates that such solution may be held over a period of twenty-four hours or greater
30 after the first use. The invention further comprises an article of manufacture, comprising packaging material, a vial comprising a formulation of FSH or an FSH variant according to the invention, wherein said packaging material comprises written material which instructs a patient to reconstitute the FSH or an FSH variant in the aqueous diluent to form a solution which may be held over a period of twenty-four
35 hours or greater.

As noted above, in a preferred embodiment, the invention provides an article of manufacture, comprising packaging material and a vial comprising freeze dried FSH or an FSH variant, LH or an LH variant, or FSH and LH, Pluronic F68. The bacteriostatic within the second container including the diluent is selected from phenol and *m*-cresol, optionally with further excipients, wherein said packaging material comprises written material which indicates that such solution may be held over a period of twenty-four hours or greater after the first use.

The range of protein hormone in the formulations of the invention includes amounts yielding upon reconstitution, concentrations from about 1.0 µg/ml to about 50 mg/ml, although lower and higher concentrations are operable and are dependent on the intended delivery vehicle, e.g., solution formulations will differ from transdermal patch, pulmonary, transmucosal, or osmotic or micro pump methods. The protein hormone concentration is preferably at or about 5.0 µg/ml to at or about 2 mg/ml, more preferably at or about 10 µg/ml to at or about 1 mg/ml, most preferably at or about 50 µg/ml to at or about 200 µg/ml.

The range of protein hormone in the formulations of the invention includes amounts yielding upon reconstitution, concentrations from about 1.0 µg/ml to about 50 mg/ml, although lower and higher concentrations are operable and are dependent on the intended delivery vehicle, e.g., solution formulations will differ from transdermal patch, pulmonary, transmucosal, or osmotic or micro pump methods. The protein hormone concentration is preferably at or about 5.0 µg/ml to at or about 2 mg/ml, more preferably at or about 10 µg/ml to at or about 1 mg/ml, most preferably at or about 50 µg/ml to at or about 200 µg/ml.

Preferably the formulations of the invention retain at least at or about 80% of the FSH activity and/or LH activity at the time of packaging over a period of 24 months (before the first use). FSH activity can be measured using the Steelman-Pohley ovarian weight gain bioassay⁵. LH activity can be measured using the rat seminal vesicle weight gain bioassay.

The liquid formulations of the present invention can be prepared by a process which comprises mixing FSH or an FSH variant, LH, or a mixture of FSH and LH and Pluronic F68 and a bacteriostatic selected from phenol and *m*-cresol as solids or dissolving FSH or an FSH variant, LH, or a mixture of FSH and LH ("protein") and

- Pluronic F68 and a bacteriostatic selected from phenol and *m*-cresol in an aqueous diluent. Mixing the components and dissolving them in an aqueous diluent is carried out using conventional dissolution and mixing procedures. To prepare a suitable formulation, for example, a measured amount of FSH or FSH variant, LH or a mixture of FSH and LH in buffered solution is combined with Pluronic F68 and a bacteriostatic selected from phenol and *m*-cresol in a buffered solution in quantities sufficient to provide the protein, Pluronic F68 and the bacteriostatic at the desired concentrations. The resulting solution is then dispensed into vials, ampoules or cartridges. Variations of this process would be recognized by one of ordinary skill in the art. For example, the order the components are added, whether additional additives are used, the temperature and pH at which the formulation is prepared, are all factors that may be optimised for the concentration and means of administration used.
- 15 In a preferred embodiment, the liquid formulations of the invention are made by preparing individual stock solutions of known concentration of all the components of the formulation (e.g. buffer sodium phosphate, sucrose, TWEEN, methionine, FSH and/or LH), and aliquoting volumetric amounts to form a "mother solution" of the same composition as the final formulation. The "mother solution" is preferably filtered through a Duropore® (Millipore) 0.22 micron PDF membrane, to remove microorganisms, and then aliquots are dispensed into individual containers, such as vials, ampoules or cartridges.
- 25 The freeze dried formulations of the present invention can be prepared by a process which comprises mixing FSH or an FSH variant, LH or an FSH variant, or a mixture of FSH and LH and Pluronic F68 as well as further excipients like an antioxidant and/or a buffer and subjecting the mixture to a lyophilisation. Mixing the components and lyophilising them is carried out using conventional procedures. To prepare a suitable formulation, for example, a measured amount of FSH or FSH variant, LH or LH variant or a mixture of FSH and LH is combined with Pluronic F68 and the resulting mixture is lyophilized and then dispensed into vials, ampoules or cartridges. Variations of this process would be recognized by one of ordinary skill in the art. For example, the order the components are added, whether additional additives are used, the temperature and pH at which the formulation is prepared, are all factors that may be optimised for the concentration and means of administration used.

The formulations of the invention can be administered using recognized devices. Examples comprising these single vial systems include pen -injector devices for delivery of a solution such as EasyJect®, Gonol -F® Pen, Humaject®, NovoPen®, B-D®Pen, AutoPen®, and OptiPen®.

5

The products presently claimed include packaging material. The packaging material provides, in addition to the information required by the regulatory agencies, the conditions under which the product may be used. The packaging material of the present invention provides instructions to the patient to reconstitute the FSH or an FSH variant in the aqueous diluent to form a solution and to use the solution over a period of twenty-four hours or greater for the two vial, wet/dry, product. For the single vial, solution product, the label indicates that such solution may be stored after first use for a period of twenty-four hours or greater, preferably for up to 12 or 14 days. The presently claimed products are useful for human pharmaceutical product use.

10
15

The stable preserved formulations may be provided to patients as clear solutions. The solution may be for single use or it may be reused multiple times and may suffice for a single or multiple cycles of patient treatment and thus provides a more convenient treatment regimen than currently available.

20

FSH or an FSH variant, LH, or mixtures of FSH and LH in either the stable or preserved formulations or solutions described herein, may be administered to a patient in accordance with the present invention via a variety of delivery methods including SC or IM injection; transdermal, pulmonary, transmucosal, implant, osmotic pump, cartridge, micro pump, oral, or other means appreciated by the skilled artisan, as well-known in the art.

25

The following examples are provided merely to further illustrate the preparation of the formulations and compositions of the invention. The scope of the invention shall not be construed as merely consisting of the following examples.

30

Example 1

Comparative formulations

Materials

Item	Manufacturer
r-hFSH Bulk used for candidate formulations	Laboratoires Serono SA
D-Mannitol (DAB, Ph Eur, BP, FU, USP, FCC, E421)	Merck
Sucrose (DAB, Ph Eur, BP, NF)	Merck
NaCl (ACS, ISO)	Merck
Na ₂ HPO ₄ 2H ₂ O (GR for analysis)	Merck
NaH ₂ PO ₄ H ₂ O (GR for analysis)	Merck
Benzyl Alcohol (GR for analysis)	Merck
m-Cresol (for synthesis)	Merck
TWEEN 20 (Polysorbate 20) (for synthesis)	Merck
Pluronic F68 (Poloxamer 188)	Sigma
L-Methionine (for biochemistry)	Merck
Ortho-phosphoric Acid 85% (Ph Eur, BP, NF)	Merck
1.5 mL glass cartridge	SFAM (siliconed at Aguetant)
Rubbers Type A	West Company
Crim caps	Aguettant
Millex-GV	Millipore
Syringe Driven Filter Unit – Durapore	
Durapore Membrane Filters 0.22 µm GV	Millipore
20 mL Plastic syringe Plastipak	Becton Dickinson
Steel Holder for filtration	Sartorius

Equipment

HPLC Systems	Detector mod. 486 or 490 Controller mod. 600S Pump mod. 626 Autosampler mod. 717	Waters	2
pH meter	Mod. 654	Metrohm	1
Osmometer	030-D	Osmomat	1

5 The following study evaluated the following parameters for a large number of formulations:

- Compatibility of surfactant and bacteriostatic
- Oxidation of alpha-subunit

10 The formulations were multi-dose formulations and contained either TWEEN 20 or Pluronic F68 as well as a bacteriostatic agent. The following three bacteriostatic agents were evaluated:

- Benzyl alcohol 0.9%
- *m*-Cresol 0.3%
- Phenol 0.5%

15

TWEEN 20 and Pluronic F68 were used at the following range of concentrations:

- TWEEN 20 : range from 10 to 100 µg/g
- Pluronic F68 : range from 10 to 100 µg/g

20 Solutions prepared are listed in Table 1.

Table 1: Comparative formulations

ID #	Na ₂ HPO ₄ 2H ₂ O (mg/g)	NaH ₂ PO ₄ H ₂ O (mg/g)	r-hFSH*	Pluronic F68 (µg/g)	TWEEN 20 (µg/g)	Bacteriostat	Excipient (mg/g)
1P	1.11	0.45	600 IU/g	10	-	0.5% Phenol	Sucrose 70.6
2P	1.11	0.45	600 IU/g	10	-	0.5% Phenol	Mannitol 38.7
3P	1.11	0.45	600 IU/g	100	-	0.5% Phenol	Sucrose 70.6
4P	1.11	0.45	600 IU/g	100	-	0.5% Phenol	Mannitol 38.7
5P	1.11	0.45	600 IU/g	-	10	0.5% Phenol	Sucrose 70.6
6P	1.11	0.45	600 IU/g	-	10	0.5% Phenol	Mannitol 38.7
7	1.11	0.45	600 IU/g	-	100	0.9% benzyl alcohol	NaCl 6.0
8	1.11	0.45	600 IU/g	-	100	0.9% benzyl alcohol	Sucrose 62.3
9	1.11	0.45	600 IU/g	-	100	0.9% benzyl alcohol	Mannitol 34.1

Table 1: Comparative formulations

ID #	Na ₂ HPO ₄ 2H ₂ O (mg/g)	NaH ₂ PO ₄ H ₂ O (mg/g)	r-hFSH*	Pluronic F68 (µg/g)	TWEEN 20 (µg/g)	Bacteriostat	Excipient (mg/g)
10	1.11	0.45	600 IU/g	-	100	0.3 % m- Cresol	NaCl 7.6
11	1.11	0.45	600 IU/g	-	100	0.3 % m- Cresol	Sucrose 78.0
12	1.11	0.45	600 IU/g	-	100	0.3 % m- Cresol	Mannitol 42.7
13	1.11	0.45	600 IU/g	-	10	0.9% benzyl alcohol	NaCl 6.0
14	1.11	0.45	600 IU/g	-	10	0.9% benzyl alcohol	Sucrose 62.3
15	1.11	0.45	600 IU/g	-	10	0.9% benzyl alcohol	Mannitol 34.1
16	1.11	0.45	600 IU/g	-	10	0.3 % m- Cresol	NaCl 7.6
17	1.11	0.45	600 IU/g	-	10	0.3 % m- Cresol	Sucrose 78.0
18	1.11	0.45	600 IU/g	-	10	0.3 % m- Cresol	Mannitol 42.7

Table 1: Comparative formulations

ID #	Na ₂ HPO ₄ 2H ₂ O (mg/g)	NaH ₂ PO ₄ H ₂ O (mg/g)	r-hFSH*	Pluronic F68 (µg/g)	TWEEN 20 (µg/g)	Bacteriostat	Excipient (mg/g)
19	1.11	0.45	600 IU/g	100	-	0.9% benzyl alcohol	NaCl 6.0
20	1.11	0.45	600 IU/g	100	-	0.9% benzyl alcohol	Sucrose 62.3
21	1.11	0.45	600 IU/g	100	-	0.9% benzyl alcohol	Mannitol 34.1
22	1.11	0.45	600 IU/g	100	-	0.3% m- Cresol	NaCl 7.6
23	1.11	0.45	600 IU/g	100	-	0.3% m- Cresol	Sucrose 78.0
24	1.11	0.45	600 IU/g	100	-	0.3% m- Cresol	Mannitol 42.7
25	1.11	0.45	600 IU/g	10	-	0.9% benzyl alcohol	NaCl 6.0
26	1.11	0.45	600 IU/g	10	-	0.9% benzyl alcohol	Sucrose 62.3
27	1.11	0.45	600 IU/g	10	-	0.9% benzyl alcohol	Mannitol 34.1

Table 1: Comparative formulations

ID #	Na ₂ HPO ₄ 2H ₂ O (mg/g)	NaH ₂ PO ₄ H ₂ O (mg/g)	r-hFSH*	Pluronic F68 (µg/g)	TWEEN 20 (µg/g)	Bacteriostat	Excipient (mg/g)
28	1.11	0.45	600 IU/g	10	-	0.3% m- Cresol	NaCl 7.6
29	1.11	0.45	600 IU/g	10	-	0.3% m- Cresol	Sucrose 78.0
30	1.11	0.45	600 IU/g	10	-	0.3% m- Cresol	Mannitol 42.7

*FSH was added to the formulations on the basis of its biopotency instead of protein content.

From visual examination of the formulations, it was determined that TWEEN 20 cannot be used with *m*-cresol and phenol because FSH formulations containing TWEEN 20 and *m*-cresol or TWEEN 20 and phenol presented a white opalescent suspension. In contrast, FSH formulations containing Pluronic F68 did not exhibit this problem with *m*-cresol and phenol. The use of Pluronic F68 permits the use of phenol and *m*-cresol.

Combination of FSH and Pluronic F68 with antioxidants

- 10 The following antioxidants were evaluated for their ability to inhibit oxidation of the α -subunit in the presence of Pluronic F68:
- Methionine : range from 10 to 100 μ g/g
 - Ascorbic Acid : range from 10 to 100 μ g/g
- 15 Sucrose and Mannitol were used as tonicity agents and TWEEN 20 or Pluronic were added at the concentration of 100 μ g/g.

The formulations prepared are listed in Table 2.

Table 2. Comparative formulations with and without methionine

ID#	Na ₂ HPO ₄ 2H ₂ O (mg/g)	NaH ₂ PO ₄ H ₂ O (mg/g)	RhFSH	Pluronic F68 (µg/g)	TWEEN (µg/g)	Ascorbic Acid (µg/g)	Methionine (µg/g)	Bacteriostat	Excipient
31	1.11	0.45	600 IU/g	100	-	-	-	0.3% m- cresol	Sucrose
32	1.11	0.45	600 IU/g	100	-	-	-	0.3% m- cresol	Mannitol
33	1.11	0.45	600 IU/g	-	100	-	-	0.9% benzyl alcohol	Sucrose
34	1.11	0.45	600 IU/g	-	100	-	-	0.9% benzyl alcohol	Mannitol
35	1.11	0.45	600 IU/g	100	-	-	-	0.9% benzyl alcohol	Sucrose
36	1.11	0.45	600 IU/g	100	-	-	-	0.9% benzyl alcohol	Mannitol
37	1.11	0.45	600 IU/g	100	-	-	10	0.3% m- cresol	Sucrose
38	1.11	0.45	600 IU/g	100	-	-	10	0.3% m- cresol	Mannitol
39	1.11	0.45	600 IU/g	100	-	-	100	0.3% m- cresol	Sucrose

Table 2. Comparative formulations with and without methionine

ID#	Na ₂ HPO ₄ 2H ₂ O (mg/g)	NaH ₂ PO ₄ H ₂ O (mg/g)	RhFSH	Pluronic F68 (µg/g)	TWEEN (µg/g)	Ascorbic Acid (µg/g)	Methionine (µg/g)	Bacteriostat	Excipient
40	1.11	0.45	600 IU/g	100	-	-	100	0.3% m- cresol	Mannitol
41	1.11	0.45	600 IU/g	100	-	10	-	0.3% m- cresol	Sucrose
42	1.11	0.45	600 IU/g	100	-	10	-	0.3% m- cresol	Mannitol
43	1.11	0.45	600 IU/g	100	-	100	-	0.3% m- cresol	Sucrose
44	1.11	0.45	600 IU/g	100	-	100	-	0.3% m- cresol	Mannitol
45	1.11	0.45	600 IU/g	-	100	-	10	0.9% benzyl alcohol	Sucrose
46	1.11	0.45	600 IU/g	-	100	-	10	0.9% benzyl alcohol	Mannitol
47	1.11	0.45	600 IU/g	-	100	-	100	0.9% benzyl alcohol	Sucrose
48	1.11	0.45	600 IU/g	-	100	-	100	0.9% benzyl alcohol	Mannitol

Table 2. Comparative formulations with and without methionine

ID#	Na ₂ HPO ₄ 2H ₂ O (mg/g)	NaH ₂ PO ₄ H ₂ O (mg/g)	RhFSH	Pluronic F68 (µg/g)	TWEEN (µg/g)	Ascorbic Acid (µg/g)	Methionine (µg/g)	Bacteriostat	Excipient
49	1.11	0.45	600 IU/g	-	100	10	-	0.9% benzyl alcohol	Sucrose
50	1.11	0.45	600 IU/g	-	100	10	-	0.9% benzyl alcohol	Mannitol
51	1.11	0.45	600 IU/g	-	100	100	-	0.9% benzyl alcohol	Sucrose
52	1.11	0.45	600 IU/g	-	100	100	-	0.9% benzyl alcohol	Mannitol
53	1.11	0.45	600 IU/g	100	-	-	10	0.9% benzyl alcohol	Sucrose
54	1.11	0.45	600 IU/g	100	-	-	10	0.9% benzyl alcohol	Mannitol
55	1.11	0.45	600 IU/g	100	-	-	100	0.9% benzyl alcohol	Sucrose
56	1.11	0.45	600 IU/g	100	-	-	100	0.9% benzyl alcohol	Mannitol
57	1.11	0.45	600 IU/g	100	-	10	-	0.9% benzyl alcohol	Sucrose

Table 2. Comparative formulations with and without methionine

ID#	Na ₂ HPO ₄ 2H ₂ O (mg/g)	NaH ₂ PO ₄ H ₂ O (mg/g)	RhFSH	Pluronic F68 (µg/g)	TWEEN (µg/g)	Ascorbic Acid (µg/g)	Methionine (µg/g)	Bacteriostat	Excipient
58	1.11	0.45	600 IU/g	100	-	10	-	0.9% benzyl alcohol	Mannitol
59	1.11	0.45	600 IU/g	100	-	100	-	0.9% benzyl alcohol	Sucrose
60	1.11	0.45	600 IU/g	100	-	100	-	0.9% benzyl alcohol	Mannitol
61	1.11	0.45	600 IU/g	100	-	-	-	Phenol	Sucrose
62	1.11	0.45	600 IU/g	100	-	-	-	Phenol	Mannitol
63	1.11	0.45	600 IU/g	-	-	-	-	Phenol	Sucrose
64	1.11	0.45	600 IU/g	100	-	-	10	Phenol	Mannitol
65	1.11	0.45	600 IU/g	100	-	-	100	Phenol	Sucrose
66	1.11	0.45	600 IU/g	100	-	-	100	Phenol	Mannitol
67	1.11	0.45	600 IU/g	100	-	10	-	Phenol	Sucrose
68	1.11	0.45	600 IU/g	100	-	10	-	Phenol	Mannitol
69	1.11	0.45	600 IU/g	100	-	100	-	Phenol	Sucrose
70	1.11	0.45	600 IU/g	100	-	100	-	Phenol	Mannitol

FSH was added to the formulations on the basis of its biopotency instead of protein content.

20 g of each formulation was prepared into Falcon polypropylene tubes and filtered through a 3cm² 0.22 µm Millex-GV Syringe Driven filter unit Durapore, then analysed for a value at t=0. The solutions were then stored at 40°C and tested according the following scheme:

Analytical test	T=0	1 week	2 weeks	3 weeks	4 weeks
Reverse Phase-HPLC for oxidised alpha subunit (%)	X	X	X	X	X
Size Exclusion-HPLC for protein quantitation (µg/g)	X	X	X	X	X
Size Exclusion-HPLC for qualitative free subunits	X	X	X	X	X

5

(X): Test performed

Reverse phase HPLC reveals that in formulations containing FSH, Pluronic F68, *m*-cresol and methionine (at 10 and 100 µg/ml), oxidation of the α-subunit of FSH when the formulation is stored at 40°C, is greatly reduced, versus a formulation containing no methionine, as can be seen in Figure 1. Based on the average of two experiments, in the Formulation containing no methionine, the percent of oxidised α-subunit is 2.3 at T=0, 4.0 at T= 1 week, and 7.1 at T= 2 weeks. In the formulation containing 10 µg/ml methionine, the percent of oxidised α-subunit is 2.0 at T=0, 3.2 at T= 1 week, and 3.8 at T= 2 weeks. In the formulation containing 100 µg/ml methionine, the percent of oxidised α-subunit is 1.8 at T=0, 1.7 at T= 1 week, and 1.3 at T= 2 weeks.

10

15

Example 2

Liquid single-dose formulation of recombinant FSH for subcutaneous or intramuscular injection

20

Based on the results of Example 1, the following formulation was prepared.

25

Components 1 to 7 listed in Table 3 were prepared as volumetric solutions in WFI. Aliquots of each solution were added to a mixing vessel to form a "mother solution". The mother solution was dispensed into vials to contain 10.9 micrograms (150 IU) or 5.45 micrograms (75 IU) of FSH.

With recombinant FSH, the bioactivity and specific activity are consistent, allowing the FSH to be filled by mass, rather than by bioassay.

Table 3. Components of FSH single dose liquid formulations			
Component #	Description	150 IU FSH	75 IU FSH
1	rhFSH ($\mu\text{g}/\text{vial}$)	10.9 (150 IU)	5.45 (75 IU)
2	Sucrose (mg/vial)	15.00	7.50
3	$\text{NaH}_2\text{PO}_4 \cdot \text{H}_2\text{O}$ (mg/vial)	0.111	0.0555
4	$\text{Na}_2\text{HPO}_4 \cdot 2\text{H}_2\text{O}$ (mg/vial)	0.273	0.1365
5	Pluronic F68 (mg/vial)	0.025	0.0125
6	Methionine (mg/vial)	0.025	0.0125
7	<i>m</i> -cresol (mg/vial)	0.75	0.375
8	PH	7.0	7.0
9	WFI	q.s. to 1 ml	q.s. to 0.5 ml

- 5 The vials were filled and sealed under sterile conditions. The formulation has a shelf life of up to two years at ambient temperatures.

Example 3

Liquid multi-dose formulation of recombinant FSH for subcutaneous or

10 Intramuscular Injection

Based on the results of Example 1, the following multi-dose formulation was prepared.

Components 1 to 7 listed in Table 4 were prepared as volumetric solutions in WFI.

- 15 Aliquots of each solution were added to a mixing vessel to form a "mother solution". The mother solution was dispensed into vials to contain 22.2 micrograms (305 IU), 33.3 micrograms (458 IU) and 66.7 micrograms (916 IU) of FSH. The resulting formulations deliver a total of 300, 450 and 900 IU of FSH.

- 20 The cartridges were filled and sealed under sterile conditions. The multi-dose formulation can be stored at at or about 2-8°C, more preferably at or about 4-5°C, until the first use for up to two years. After the first use, the cartridge should be

stored at at or about 2-8°C, more preferably at or about 4-5°C, over the multi-dose period, which may be 24 hours, 2 days, or up to 12 or 14 days.

Table 4. Components of FSH multi-dose liquid formulations				
Component #	Description	300 IU FSH	450 IU FSH	900 IU FSH
1	rhFSH (µg/cartridge)	22.2 (305 IU)	33.3 (458 IU)	66.7 (916 IU)
2	Sucrose (mg/cartridge)	30.0	45.0	90.0
3	NaH ₂ PO ₄ ·H ₂ O (mg/cartridge)	0.225	0.337	0.675
4	Na ₂ HPO ₄ ·2H ₂ O (mg/cartridge)	0.555	0.832	1.665
5	Pluronic F68 (mg/vial)	0.050	0.075	0.150
6	Methionine (mg/vial)	0.050	0.075	0.150
7	<i>m</i> -cresol (mg/vial)	1.50	2.25	4.50
8	pH	7.0	7.0	7.0
9	WFI	q.s. to 0.5 ml	q.s. to 0.75 ml	q.s. to 1.5 ml

5 Example 4

Liquid single-dose formulation of recombinant LH for subcutaneous or intramuscular injection

The following formulation was prepared.

- 10 Components 1 to 7 listed in Table 5 were prepared as volumetric solutions in WFI. Aliquots of each solution were added to a mixing vessel to form a "mother solution". The mother solution was dispensed into vials to contain 3 micrograms (75 IU) of LH. The resulting formulation delivers a single dose of 75 IU LH.
- 15 With recombinant LH, the bioactivity and specific activity are consistent, allowing the LH to be filled by mass, rather than by bioassay.

Table 5. Components of LH single dose liquid formulation		
Component #	Description	LH 75 IU
1	rhLH ($\mu\text{g}/\text{vial}$)	3.0
2	Sucrose (mg/vial)	52.5
3	$\text{NaH}_2\text{PO}_4 \cdot \text{H}_2\text{O}$ (mg/vial)	0.052
4	$\text{Na}_2\text{HPO}_4 \cdot 2\text{H}_2\text{O}$ (mg/vial)	0.825
5	Pluronic F68 (mg/vial)	0.0125
6	Methionine (mg/vial)	0.125
7	<i>m</i> -cresol (mg/vial)	0.375
9	WFI	q.s. to 0.5 ml

The vials were filled and sealed under sterile conditions. The formulation has a shelf life of up to two years.

5

Example 5

Liquid multi-dose formulations of recombinant FSH and LH (2:1) for subcutaneous or intramuscular injection

The following multi-dose formulations of FSH and LH were prepared, with FSH:LH ratio of 2:1.

10

Components 1 to 8 listed in Table 6 were prepared as volumetric solutions in WFI. Aliquots of each solution were added to a mixing vessel and mixed to form a "mother solution". The pH of the mother solution was adjusted to 8.0, if necessary, by addition of NaOH or HCl. The mother solution was dispensed into cartridges to contain 18.3 micrograms LH (457 IU) with 66.7 micrograms FSH (916 IU), intended for 6 doses of 150 IU FSH each; 9.2 micrograms LH (230 IU) with 33.3 micrograms FSH (458 IU), intended for 3 doses of 150 IU FSH each; and 6.1 micrograms LH (152.5 IU) with 22.23 micrograms FSH (305 IU), intended for 2 doses of 150 IU FSH each.

15

20

The cartridges were filled and sealed under sterile conditions. The multi-dose formulation can be stored at at or about 2-8°C, more preferably at or about 4-5°C until

the first use for up to two years. After the first use, the cartridge should be stored at at or about 2-8°C, more preferably at or about 4-5°C, over the multi-dose period, which may be 24 hours, 2 days, or up to 12 or 14 days.

Table 6. Components of FSH and LH (2:1) multi-dose liquid formulations				
Component #	Description	6 doses	3 doses	2 doses
1	rhLH (µg/cartridge)	18.3 (457 IU)	9.2 (230 IU)	6.1 (152.5 IU)
2	rhFSH (µg/cartridge)	66.7 (916 IU)	33.3 (458 IU)	22.23 (305 IU)
3	Sucrose (mg/cartridge)	115.5	57.75	38.5
4	H ₃ PO ₄ (mg/cartridge)	1.35	0.735	0.49
5	NaOH (mg/cartridge)	q.s. to pH 8.0	q.s. to pH 8.0	q.s. to pH 8.0
6	Pluronic F68 (mg/vial)	375.0	187.5	125.0
7	Methionine (µg/cartridge)	225	112.5	75.0
8	m-cresol (mg/cartridge)	4.5	2.25	1.5
9	pH	8.0	8.0	8.0
10	WFI	q.s. to 1.5 ml	q.s. to 0.75 ml	q.s. to 0.5 ml

5

Example 6

Liquid multi-dose formulations of recombinant FSH and LH (1:1) for subcutaneous or intramuscular injection

10 The following multi-dose formulations of FSH and LH were prepared, with FSH:LH ratio of 1:1.

Components 1 to 8 listed in Table 7 were prepared as volumetric solutions in WFI. Aliquots of each solution were added to a mixing vessel and mixed to form a "mother solution". The pH of the mother solution was adjusted to 8.0, if necessary, by addition of NaOH or HCl. The mother solution was dispensed into cartridges to contain 36.6 micrograms LH (914 IU) with 66.7 micrograms FSH (916 IU), intended

15

for 6 doses of 150 IU FSH each; 18.4 micrograms LH (460 IU) with 33.3 micrograms FSH (458 IU), intended for 3 doses of 150 IU FSH each; and 12.2 micrograms LH (305 IU) with 22.23 micrograms FSH (305 IU), intended for 2 doses of 150 IU FSH each.

5

The cartridges were filled and sealed under sterile conditions. The multi-dose formulation can be stored at or about 2-8°C, more preferably at or about 4-5°C until the first use for up to two years. After the first use, the cartridge should be stored at or about 2-8°C, more preferably at or about 2-8°C, more preferably at or about 4-5°C, over the multi-dose period, which may be 24 hours, 2 days, or up to 12 or 14 days.

10

Component #	Description	6 doses	3 doses	2 doses
1	rhLH (µg/cartridge)	36.6 (914 IU)	18.4 (460 IU)	12.2 (305 IU)
2	rhFSH (µg/cartridge)	66.7 (916 IU)	33.3 (458 IU)	22.23 (305 IU)
3	Sucrose (mg/cartridge)	115.5	57.75	38.5
4	H ₃ PO ₄ (mg/cartridge)	1.35	0.735	0.49
5	NaOH (mg/cartridge)	q.s. to pH 8.0	q.s. to pH 8.0	q.s. to pH 8.0
6	Pluronic F68 (mg/vial)	375.0	187.5	125.0
7	Methionine (µg/cartridge)	225	112.5	75.0
8	m-cresol (mg/cartridge)	4.5	2.25	1.5
9	pH	8.0	8.0	8.0
10	WFI	q.s. to 1.5 ml	q.s. to 0.75 ml	q.s. to 0.5 ml

Example 7

Stability experiments for liquid multi-dose formulations of FSH mixed with LH

7.1. Reverse phase HPLC analysis for protein content

The formulation of Example 5 (6 doses) was evaluated for protein content for both
5 FSH and LH, using a reverse-phase HPLC method.

Protein content (FSH and LH) was measured at zero time, and after 1, 2, 3 and 6
months storage of the formulation at 4°C. The results are listed in Table 8 as
micrograms of FSH or LH per gram of solvent.

10

7.2. Assay of oxidised alpha-subunit

The percentage of oxidised alpha-subunit in a formulation of Example 5 was
measured by a reverse phase HPLC (RP-HPLC) method.

15 The percentage of oxidised alpha-subunit was measured at zero time, and after 1, 2,
3 and 6 months storage at 4°C. The results are listed in Table 8.

7.3. In vivo assay for FSH

The formulation of Example 5 (6 doses) was tested for FSH activity using the
20 Steelman-Pohley ovarian weight gain bioassay at zero time, and after 1, 2, 3 and 6
months of storage at 4°C. The results are listed in Table 8 as international units (IU)
per gram of solvent.

7.4. In vivo assay for LH

25 The formulation of Example 5 (6 doses) was tested for LH activity using the rat
seminal vesicle weight gain bioassay at zero time, and after 1, 2, 3 and 6 months of
storage at 4°C. The results are listed in Table 8 as international units (IU) per gram
of solvent.

7.5. Evaluation of free subunit (rFSH + rLH)

30 For a formulation of Example 5 the percentage of free subunit was evaluated by
SDS-PAGE.

Measurements were made at zero time, and after 1, 2, 3 and 6 months storage at
35 4°C. The results are reported as a percentage of the total protein (rFSH + rLH), and
are listed in Table 8.

7.5. Evaluation of aggregates

For a formulation of Example 5, the percentage of aggregates was evaluated by SDS-PAGE as described above for evaluation of free subunit in 7.5., except that higher molecular weight aggregates were determined as a percentage of the total protein (rFSH + rLH). Measurements were made at zero time and after 1, 2, 3 and 6 months storage at 4°C. Results are listed in Table 8.

7.6. Visible particles

The formulation of Example 5 was evaluated visually for particles at zero time, and after 3 and 6 months of storage at 4°C. Results are reported in Table 8.

7.7. pH

The pH of a formulation of Example 5 was measured at zero time and after 1, 2, 3 and 6 months storage at 4°C. Results are listed in Table 8.

Table 8. Analytical parameters for a liquid formulation of FSH and LH (2:1) at zero time and after storage at 4°C for 1, 2, 3 and 6 months					
Assay	Zero time	1 month	2 months	3 months	6 months
rFSH content by RP-HPLC (micrograms/g)	46.50	46.98	46.71	46.31	44.98
rLH content by RP-HPLC (micrograms/g)	11.74	11.81	12.68	12.67	13.21
% alpha-subunit oxidised	2.29	2.17	2.08	2.48	2.95
In vivo assay for FSH 553(IU/g)	566	Not tested	Not tested	Not tested	578 (23 weeks)
In vivo assay for LH (IU/g)	331	Not tested	Not tested	311	286
SDS-PAGE free subunit (rFSH + rLH; %)	≤ 5	Not tested	Not tested	≤ 5	≤ 5 (23 weeks)
SDS-PAGE aggregates (rFSH + rLH; %)	≤ 2	Not tested	Not tested	> 3	4
Visible particles	Free	Not tested	Not tested	Free	Free
pH	8.262	8.215	8.216	8.188	8.283

Example 8**FSH and LH freeze dried multidose formulation**

Two freeze dried formulations A and B having the following compositions have been prepared:

5	Formulation A	
	FSH	μg 32.75 (450 I.U.)
	LH	μg 9.0 (225 I.U.)
	Sucrose	mg 15.0
	$\text{NaH}_2\text{PO}_4 \cdot \text{H}_2\text{O}$	mg 0.052
10	$\text{Na}_2\text{HPO}_4 \cdot 2\text{H}_2\text{O}$	mg 0.825
	Pluronic F-68	mg 0.05
	L-Methionine	mg 0.05
	Formulation B	
	FSH	μg 65.5 (900 I.U.)
15	LH	μg 18.0 (450 I.U.)
	Sucrose	mg 30.0
	$\text{NaH}_2\text{PO}_4 \cdot \text{H}_2\text{O}$	mg 0.104
	$\text{Na}_2\text{HPO}_4 \cdot 2\text{H}_2\text{O}$	mg 1.65
	Pluronic F68	mg 0.10
20	L-Methionine	mg 0.10

The manufacturing process consists in mixing the drug substance directly with the ingredients, filtrating the solution obtained and lyophilising the filtrated.

A description of each step of the process is given in the following :

- 25 - add in a tared container WFI, di-sodium hydrogen phosphate dihydrate, sodium dihydrogen phosphate monohydrate, Sucrose, Pluronic F68 at 5% and L-methionine and stir for 10 minutes until complete dissolution.
- check the pH and eventually correct it to $\text{pH } 7.00 \pm 0.2$ with NaOH 10% or diluted H_3PO_4 .
- 30 - add FSH and LH to the above prepared mixture and gently stir the solution obtained for 10 minutes.
- check the pH again and eventually adjust it to 7.0 ± 0.1 with 10% NaOH or diluted H_3PO_4 .

- filter the solution with a 0.22 μm Durapore membrane with a filtration ratio not less than 15g/cm², under Nitrogen gas flow with a pressure not higher than 1.5 atm.
- collect the solution in a previously sterilised flask.
- 5 - fill the filtered solution into the glass container, seat the stopper and place the filled vials into a stainless steel tray.
- load the trays into the freeze dryer and lyophilise the product using the following freeze drying cycle :
 - equilibrate at +4°C for about 20 mins.
 - 10 • bring the shelves temperature at -25°C and maintain for 2 hours.
 - bring the shelves temperature at -15°C and maintain for 1 hour.
 - bring the shelves temperature at -45°C and maintain for 3 hours.
 - bring condenser temperature at -65°C.
 - apply vacuum to the chamber.
 - 15 • When the vacuum reaches a value of 7×10^{-2} mBar raise shelf temperature up to -10°C and maintain for 14 hours.
 - raise the shelf temperature up to +35°C in 8 hours and maintain up to the end of the cycle (14 hours).
 - break the vacuum allowing dry nitrogen into the chamber.
 - 20 • perform the stoppering by automatic system of the freeze dryer.
 - seal the stoppered vials with the appropriate flip-off caps.

The formulations A and B have been stored at $25 \pm 2^\circ\text{C}$, and tested for stability and biological activity as pointed out below. Prior to analysing the compositions, they are reconstituted using water for Injection comprising 0.3% of m-Cresol as bacteriostatic agent.

The stability and biological activity values were determined as follows :

- In vivo assay for FSH : The formulation was tested for FSH activity using the Steelman-Pohley ovarian weight gain bioassay
- In vivo assay for LH : The formulation was tested for LH activity using the rat seminal vesicle weight gain bioassay.
- 30 • Assay of oxidised alpha-subunit : The percentage of oxidised alpha-subunit was measured by a reverse phase HPLC (RP-HPLC) method.
- Evaluation of free subunit (rFSH + rLH) : The percentage of free subunit was evaluated by SDS-PAGE.

- Evaluation of aggregates : The percentage of aggregates was evaluated by SDS-PAGE as described above for evaluation of free subunit.

The biological tests have been performed in compliance with the regulations of the European Pharmacopeia. In particular the tests are reported in the "Menotropin" monography.

Table 9 summarizes the results of the analytical tests related to stability and biological activity of formulation A. The values were determined at 4 check-points : at time zero, after 1 month, 3 months and 6 months of storage, at a storage temperature of $25 \pm 2^\circ\text{C}$.

TABLE 9

TEST	TIME ZERO	1 MONTHS	3 MONTHS	6 MONTHS
Biological activity I.U. FSH	416	420	415	417
Biological activity I.U. LH	276	250	259	270
% oxidised product	1.95	1.81	1.95	1.57
% dimers/aggregates	<2	<2	<2	<2
% free subunits	<5	<5	<5	<5

Table 10 summarizes the results of the analytical tests related to stability and biological activity of formulation B. The values were determined at 4 check-points : at time zero, after 3 month, 6 months and 9 months of storage, at a storage temperature of $25 \pm 2^\circ\text{C}$.

TABLE 10

TEST	TIME ZERO	3 MONTHS	6 MONTHS	9 MONTHS
Biological activity I.U. FSH	821	850	830	838
Biological activity I.U. LH	570	564	580	622
% oxidised product	1.0	0.9	1.0	1.0
% dimers/aggregates	<2	<2	<2	<2
% free subunits	<5	<5	<5	<5

From TABLE 9 and 10 it may be concluded that the biological activity of formulations A and B is well conserved after 9 months of storage. The formulations have a high stability.

- The high stability is not affected by large amounts of recombinant FSH and recombinant LH.

49

Sequences:

SEQ ID NO. 1: human glycoprotein α -subunit;

SEQ ID NO. 2: hFSH β -subunit

SEQ ID NO. 3: hFSH β -subunit variant 1

5 SEQ ID NO. 4: hFSH β -subunit variant 2

SEQ ID NO. 5: hFSH β -subunit variant 3

SEQ ID NO. 6: hLH β -subunit

10

References

- ¹ Burgues et al.; *Subcutaneous self-administration of highly purified follicle stimulating hormone and human chorionic gonadotrophin for the treatment of male hypogonadotrophic hypogonadism. Spanish Collaborative Group on Male Hypogonadotrophic Hypogonadism ; Hum. Reprod.*; 1997, 12, 980-6;
- ² Shome et al., *J. Clin. Endocrinol. Metab.* 39:187-205 (1974); Shome, et al., *J. Prot. Chem.* 7:325-339, 1988;
- ³ Keutmann et al.; *Structure of human luteinizing hormone beta subunit: evidence for related carboxyl-terminal sequence among certain peptide hormones ; Biochem. Biophys. Res. Commun.*; 1979, 90, 842-848; Talmadge et al. ; *Evolution of the genes for the beta subunits of human chorionic gonadotropin and luteinizing hormone ; Nature*; 1984, 307, 37-40; Fiddes & Talmadge; *Structure, expression, and evolution of the genes for the human glycoprotein hormones; Recent Prog. Horm. Res.* ; 1984, 40, 43-78
- ⁴ Reichert LE, Ramsey RB; *Dissociation of human follicle -stimulating hormone; J. Biol. Chem.*; 1975, 250, 3034-3040
- ⁵ Klein et al.; *Pharmacokinetics and pharmacodynamics of single -chain recombinant human follicle-stimulating hormone containing the human chorionic gonadotrophin carboxyterminal peptide in the rhesus monkey ; Fertility & Sterility*; 2002, 77, 1248-1255
- ⁶ a) Fiddes, J.C., et al., *J of Mol. and Applied Genetics*, 1:3 -18(1981); b) Esch F.S., et al. *DNA* 5:363-369(1986); c) Watkins P.C., et al., *DNA* 6:205-212(1987); d) Hirai T., et al., *J. Mol. Endocrinol.* 5:147-158(1990); e) Maurer, R.A., et al., *Mol. Endocrinol.* 1:717 -723(1987); f) Guzman K., et al., *DNA Cell Biol.* 10:593 -601(1991); g) Kumar TR, et al., *Gene*. 1995 Dec 12;166(2):335-6; h) Kumar TR, et al., *Gene*. 1995 Dec 12;166(2):333 -4
- ⁷ *Biochem. Biophys. Res. Commun.*; 1979, 90, 842-848
- ⁸ Steelman et al.; *Assay of the follicle stimulating hormone based on the augmentation with human chorionic gonadotrophin; Endocrinology*; 1953, 53, 604 -616
- ⁹ Van Hell et al.; *Effects of human menopausal gonadotrophin preparations in different bioassay methods ; Acta Endocrinologica*; 1964, 47, 409-418
- ¹⁰ Van Hell et al.; *Effects of human menopausal gonadotrophin preparations in different bioassay methods ; Acta Endocrinologica*; 1964, 47, 409-418

Claims

1. A liquid pharmaceutical composition comprising follicle-stimulating hormone (FSH) or a variant thereof, as well as a surfactant selected from Pluronic® F77,
5 Pluronic F87, Pluronic F88 and Pluronic F68.
2. A liquid pharmaceutical composition comprising follicle-stimulating hormone (FSH) or a variant and luteinising hormone (LH) or a variant thereof, as well as a surfactant selected from Pluronic® F77, Pluronic F87, Pluronic F88 and
10 Pluronic F68.
3. A liquid pharmaceutical composition comprising luteinising hormone (LH) or a variant thereof, as well as a surfactant selected from Pluronic® F77, Pluronic F87, Pluronic F88 and Pluronic F68.
- 15 4. A liquid pharmaceutical composition according to any of the preceding claims, wherein the follicle-stimulating hormone (FSH) is present at a concentration of at or about 150 IU/ml to at or about 1'200 IU/ml.
- 20 5. A liquid pharmaceutical composition according to claim 4, wherein the follicle-stimulating hormone (FSH) is present at a concentration of at or about 300 IU/ml to at or about 900 IU/ml.
- 25 6. A liquid pharmaceutical composition according to claim 5, wherein the follicle-stimulating hormone (FSH) is present at a concentration of at or about 600 IU/ml.
7. A liquid pharmaceutical composition according to any of claims 2 or 3, wherein the luteinising hormone (LH) is present at a concentration of at or about 150 IU/ml to at or about 1'200 IU/ml.
30
8. A liquid pharmaceutical composition according to claim 7, wherein the luteinising hormone (LH) is present at a concentration of at or about 300 IU/ml to at or about 750 IU/ml.


9. A freeze-dried formulation comprising follicle-stimulating hormone (FSH) or a variant thereof, as well as a surfactant selected from Pluronic® F77, Pluronic F87, Pluronic F88 and Pluronic F68.
- 5 10. A freeze-dried formulation comprising luteinising hormone (LH) or a variant thereof, as well as a surfactant selected from Pluronic® F77, Pluronic F87, Pluronic F88 and Pluronic F68.
- 10 11. A freeze-dried formulation comprising follicle-stimulating hormone (FSH) or a variant thereof as well as luteinising hormone (LH) or a variant thereof, as well as a surfactant selected from Pluronic® F77, Pluronic F87, Pluronic F88 and Pluronic F68.
12. The freeze dried formulation according to any of claims 9 to 11, wherein the follicle-stimulating hormone (FSH) is present at a concentration (w/w) of at or about 0.1 to 10 µg/mg of the total formulation.
- 15 13. The freeze dried formulation according to claim 12, wherein the follicle-stimulating hormone (FSH) is present at a concentration of at or about 0.3 to 5 µg/mg of the total formulation.
14. The freeze dried formulation according to claim 13, wherein the follicle-stimulating hormone (FSH) is present at a concentration of at or about 0.37 to 2 µg/mg of the total formulation.
- 20 15. The freeze dried formulation according to any of claims 9 to 11, wherein the luteinising hormone (LH) is present at a concentration of at or about 0.1 to 3 µg/mg of the total formulation.
- 25 16. The freeze dried formulation according to claim 15, wherein the luteinising hormone (LH) is present at a concentration of at or about 0.1 to 1 µg/mg of the total formulation.
17. The freeze dried formulation according to claim 16, wherein the luteinising hormone (LH) is present at a concentration of at or about 0.1 to 0.6 µg/mg of the total formulation.
- 30 18. A pharmaceutical composition according to any of claim 1 to 17, wherein the surfactant is Pluronic F68.

19. A pharmaceutical composition according to any of the preceding claims,
wherein the follicle-stimulating hormone is human follicle-stimulating hormone
and/or the luteinising hormone (LH) is human luteinising hormone (LH).
- 5
20. A pharmaceutical composition according to claim 19, wherein the
follicle-stimulating hormone is urinary human follicle-stimulating hormone
and/or the luteinising hormone (LH) is urinary human luteinising hormone (LH).
- 10
21. A pharmaceutical composition according to claim 19, wherein the
follicle-stimulating hormone is recombinant human follicle-stimulating hormone
and/or the luteinising hormone (LH) is recombinant human luteinising hormone
(LH).
- 15
22. A pharmaceutical composition according to any of the preceding claims,
wherein the ratio of FSH to LH is within the range of at or about 6:1 to at or
about 1:6.
- 20
23. A pharmaceutical composition according to claim 22, wherein the ratio of FSH
to LH is within the range of at or about 4:1 to at or about 1:2.
24. A pharmaceutical composition according to claim 23, wherein the ratio of FSH
to LH is within the range of at or about 3:1 to at or about 1:1.
- 25
25. A pharmaceutical composition according to claim 24, wherein the ratio of FSH
to LH is within the range of at or about 2:1 and 1:1.
26. A pharmaceutical composition according to any of the preceding claims, further
comprising a bacteriostatic agent selected from phenol and *m*-cresol.
- 30
27. A pharmaceutical composition according to claim 26, wherein the bacteriostatic
agent is *m*-cresol.
28. A pharmaceutical composition according to claim 27, comprising *m*-cresol at a
concentration of at or about 0.3% (mass/mass solvent).
- 35

29. A pharmaceutical composition according to any of the preceding claims, further comprising sucrose.
30. A pharmaceutical composition according to any of the preceding claims, further comprising methionine.
31. A pharmaceutical composition according to any of the preceding claims, further comprising a phosphate buffer at a pH of at or about 6.0 to at or about 8.0.
32. A pharmaceutical composition according to claim 31, further comprising a phosphate buffer at a pH of at or about 7.0.
33. A pharmaceutical composition according to claim 32, comprising the following ingredients: rFSH, Pluronic F68, sucrose, methionine, *m*-cresol, and an aqueous phosphate buffer at a pH of at or about 7.0.
34. A pharmaceutical composition according to claim 33, wherein the rFSH is present at a concentration of at or about 600 IU/ml, the Pluronic F68 is present at a concentration of at or about 0.1 mg/ml, the sucrose is present at a concentration of at or about 60 mg/ml, the methionine is present at a concentration of at or about 0.1 mg/ml, the *m*-cresol is present at a concentration of at or about 3 mg/ml, and the phosphate buffer is at or about 10 mM in phosphate.
35. A freeze dried formulation according to claim 11, comprising 32.75 µg of recombinant FSH, 9.0 µg of recombinant LH, 15.0 mg of sucrose, 0.052 mg of NaH₂PO₄ H₂O, 0.825 mg of Na₂HPO₄ 2H₂O, 0.05 mg of Pluronic F68 and 0.05 mg of L-methionine.
36. The freeze dried formulation according to claim 11, comprising 65.5 µg of recombinant FSH, 18.0 µg of recombinant LH, 30.0 mg of sucrose, 0.104 mg of NaH₂PO₄ H₂O, 1.65 mg of Na₂HPO₄ 2H₂O, 0.10 mg of Pluronic F68 and 0.10 mg of L-methionine.
37. A method for manufacturing a pharmaceutical composition comprising the step of forming a solution of FSH, and a surfactant selected from Pluronic® F77, Pluronic F87, Pluronic F88 and Pluronic F68, and a liquid diluent.

38. A method according to claim 37, wherein the surfactant is Pluronic F68.
39. A method according to claim 37 or 38, comprising the further step of adding a bacteriostatic agent selected from phenol and *m*-cresol.
- 5 40. A method for manufacturing a packaged pharmaceutical composition comprising placing a solution comprising FSH, and a surfactant selected from Pluronic® F77, Pluronic F87, Pluronic F88 and Pluronic F68, in a vial, ampoule or cartridge.
- 10 41. A method according to claim 40, wherein the surfactant is Pluronic F68.
42. An article of manufacture comprising a first container filled with a freeze dried formulation according to any claims from 9 to 11 and a second container comprising a solvent for reconstitution.
- 15 43. An article of manufacture according to claim 42, whereby the second container comprises an aqueous diluent containing *m*-cresol.
- 20 44. A method for manufacturing a freeze dried formulation according to any of claims 9 to 11, comprising the step of forming a mixture of FSH with or without LH, or LH alone LH as well as a surfactant selected from Pluronic® F77, Pluronic F87, Pluronic F88 and Pluronic F68, and subjecting the mixture to a lyophilisation.
45. A method according to claim 44, wherein the surfactant is Pluronic F68.

Figure 1



SEQUENCE LISTING

<110> ARES TRADING SA

<120> FSH and FSH variant formulations

<130> US 847 Y

<160> 6

<170> PatentIn version 3.1

<210> 1

<211> 91

<212> PRT

<213> Homo sapiens

<400> 1

Ala Pro Asp Val Gln Asp Cys Pro Glu Cys Thr Leu Gln Glu Asn Pro
1 5 10 15

Phe Phe Ser Gln Pro Gly Ala Pro Ile Leu Gln Cys Met Gly Cys Cys
20 25 30

Phe Ser Arg Ala Tyr Pro Thr Pro Leu Arg Ser Lys Lys Thr Met Leu
35 40 45

Val Gln Lys Asn Val Thr Ser Glu Ser Thr Cys Cys Val Ala Lys Ser
50 55 60

Tyr Asn Arg Val Thr Val Met Gly Gly Phe Val Glu Asn His Thr Ala
65 70 75 80

Cys His Cys Ser Thr Cys Tyr Tyr His Lys Ser
85 90

<210> 2

<211> 129

<212> PRT

<213> Homo sapiens

<400> 2

Met Lys Thr Leu Gln Phe Phe Phe Leu Phe Cys Cys Trp Lys Ala Ile
 1 5 10 15
 Cys Cys Asn Ser Cys Glu Leu Thr Asn Ile Thr Ile Ala Ile Glu Lys
 20 25 30
 Glu Glu Cys Arg Phe Cys Ile Ser Ile Asn Thr Thr Trp Cys Ala Gly
 35 40 45
 Tyr Cys Tyr Thr Arg Asp Leu Val Tyr Lys Asp Pro Ala Arg Pro Lys
 50 55 60
 Ile Gln Lys Thr Cys Thr Phe Lys Glu Leu Val Tyr Glu Thr Val Arg
 65 70 75 80
 Val Pro Gly Cys Ala His His Ala Asp Ser Leu Tyr Thr Tyr Pro Val
 85 90 95
 Ala Thr Gln Cys His Cys Gly Lys Cys Asp Ser Asp Ser Thr Asp Cys
 100 105 110
 Thr Val Arg Gly Leu Gly Pro Ser Tyr Cys Ser Phe Gly Glu Me t Lys
 115 120 125

Glu

<210> 3

<211> 108

<212> PRT

<213> Homo sapiens

<400> 3

Asn Ser Cys Glu Leu Thr Asn Ile Thr Ile Ala Ile Glu Lys Glu Glu

```

1              5              10              15
Cys Arg Phe Cys Ile Ser Ile Asn Thr Thr Trp Cys Ala Gly Tyr Cys
      20              25              30
Tyr Thr Arg Asp Leu Val Tyr Lys Asp Pro Ala Arg Pro Lys Ile Gln
      35              40              45
Lys Thr Cys Thr Phe Lys Glu Leu Val Tyr Glu Thr Val Arg Val Pro
      50              55              60
Gly Cys Ala His His Ala Asp Ser Leu Tyr Thr Tyr Pro Val Ala Thr
65              70              75              80
Gln Cys His Cys Gly Lys Cys Asp Ser Asp Ser Thr Asp Cys Thr Val
      85              90              95
Arg Gly Leu Gly Pro Ser Tyr Cys Ser Phe Gly Glu
      100              105

```

```
<210> 4
<211> 106
<212> PRT
<213> Homo sapiens
```

<400> 4

Asn Ser Cys Glu Leu Thr Asn Ile Ala Ile Glu Lys Glu Glu Cys Arg
 1 5 10 15
 Phe Cys Ile Ser Ile Asn Thr Trp Cys Ala Gly Tyr Cys Tyr Thr Arg
 20 25 30
 Asp Leu Val Tyr Lys Asp Pro Ala Arg Pro Lys Ile Gln Lys Thr Cys
 35 40 45
 Thr Phe Lys Glu Leu Val Tyr Glu Thr Val Arg Val Pro Gly Cys Ala
 50 55 60
 His His Ala Asp Ser Leu Tyr Thr Val Pro Val Ala Thr Gln Cys His
 65 70 75 80
 Cys Gly Lys Cys Asp Ser Asp Ser Thr Asp Cys Thr Val Arg Gly Leu
 85 90 95

Gly Pro Ser Tyr Cys Ser Phe Gly Glu Met
100 105

<210> 5

<211> 110

<212> PRT

<213> Homo sapiens

<400> 5

Asn Ser Cys Glu Leu Thr Asn Ile Thr Ile Ala Ile Glu Lys Glu Glu
1 5 10 15

Cys Arg Phe Cys Ile Ser Ile Asn Thr Thr Trp Cys Ala Gly Tyr Cys
20 25 30

Tyr Thr Arg Asp Leu Val Tyr Lys Asp Pro Ala Arg Pro Lys Ile Gln
35 40 45

Lys Thr Cys Thr Phe Lys Glu Leu Val Tyr Glu Thr Val Arg Val Pro
50 55 60

Gly Cys Ala His His Ala Asp Ser Leu Tyr Thr Tyr Pro Val Ala Thr
65 70 75 80

Gln Cys His Cys Gly Lys Cys Asp Ser Asp Ser Thr Asp Cys Thr Val
85 90 95

Arg Gly Leu Gly Pro Ser Tyr Cys Ser Phe Gly Glu Met Lys
100 105 110

<210> 6

<211> 112

<212> PRT

<213> Homo sapiens

<400> 6

Ser Arg Glu Pro Leu Arg Pro Trp Cys His Pro Ile Asn Ala Ile Leu
1 5 10 15

Ala Val Glu Lys Glu Gly Cys Pro Val Cys Ile Thr Val Asn Thr Thr
20 25 30

Ile Cys Ala Gly Tyr Cys Pro Thr Met Arg Val Leu Gln Ala Val Leu
35 40 45

Pro Pro Leu Pro Gln Val Cys Thr Tyr Arg Asp Val Arg Phe Glu Ser
50 55 60

Ile Arg Leu Pro Gly Cys Pro Arg Gly Val Asp Pro Val Val Ser Phe
65 70 75 80

Pro Val Ala Leu Ser Cys Arg Cys Gly Pro Cys Arg Arg Ser Thr Ser
85 90 95

Asp Cys Gly Gly Pro Lys Asp His Pro Leu Thr Cys Asp His Pro Gln
100 105 110

Box No. I Nucleotide and/or amino acid sequence(s) (Continuation of item 1.b of the first sheet)

1. With regard to any nucleotide and/or amino acid sequence disclosed in the international application and necessary to the claimed invention, the international search was carried out on the basis of:
- a. type of material
- ☒ a sequence listing
- ☐ table(s) related to the sequence listing
- b. format of material
- ☒ in written format
- ☒ in computer readable form
- c. time of filing/furnishing
- ☒ contained in the international application as filed
- ☐ filed together with the international application in computer readable form
- ☒ furnished subsequently to this Authority for the purpose of search
2. ☒ In addition, in the case that more than one version or copy of a sequence listing and/or table relating thereto has been filed or furnished, the required statements that the information in the subsequent or additional copies is identical to that in the application as filed or does not go beyond the application as filed, as appropriate, were furnished.
3. Additional comments:

INTERNATIONAL SEARCH REPORT

International application No.

PCT/EP2004/050432

Box No. IV Text of the abstract (Continuation of Item 5 of the first sheet)

The invention relates to the field of pharmaceutical formulations of follicle-stimulating hormone (FSH), luteinising hormone (LH), and mixtures of FSH and luteinising hormone (LH), and to methods of producing such formulations.

The invention provides a liquid or freeze-dried formulation of FSH, or LH, or FSH and LH comprising a surfactant selected from Pluronic® F77, Pluronic F87, Pluronic F88 and Pluronic F68.

INTERNATIONAL SEARCH REPORT

International Application No
T/EP2004/050432A. CLASSIFICATION OF SUBJECT MATTER
IPC 7 A61K47/10 A61K38/24

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
IPC 7 A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, PAJ, EMBASE, MEDLINE, CHEM ABS Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	EP 0 974 359 A (LILLY CO ELI) 26 January 2000 (2000-01-26) cited in the application paragraphs '0014! - '0020! paragraph '0025! paragraphs '0074! - '0076! paragraph '0081!; example 12 paragraph '0083! paragraph '0125!	1-45
Y	EP 0 853 945 A (AKZO NOBEL NV) 22 July 1998 (1998-07-22) cited in the application page 3, lines 51,52 page 4, lines 23-37	1-45

☒ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
- *&* document member of the same patent family

Date of the actual completion of the international search

25 August 2004

Date of mailing of the international search report

13/09/2004

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Deck, A

INTERNATIONAL SEARCH REPORT

International Application No
T/EP2004/050432

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	US 5 811 096 A (ALEMAN CLAUDE ET AL) 22 September 1998 (1998-09-22) column 1, lines 47-52 column 2, lines 30-51 column 3, lines 5-42	1-45
Y	US 6 066 620 A (CHANG C PAUL ET AL) 23 May 2000 (2000-05-23) column 1, lines 26-60 column 3, lines 53-55 column 6, lines 18-24 column 16, lines 25-34; claims 1-3	1-45
Y	EP 0 318 081 A (AKZO NV) 31 May 1989 (1989-05-31) examples	1-45
Y	FR 2 782 455 A (SANOFI SA) 25 February 2000 (2000-02-25) page 2, lines 1-25 page 4, lines 3-14; claim 10	1-45
Y	EP 0 603 159 A (GENENTECH INC) 22 June 1994 (1994-06-22) claims	1-45
Y	EP 0 448 146 A (AKZO NV) 25 September 1991 (1991-09-25) cited in the application page 4, line 31 - page 5, line 27; examples	1-45
Y	US 5 650 390 A (SAMARITANI FABRIZIO ET AL) 22 July 1997 (1997-07-22) cited in the application the whole document.	1-45
A	US 5 356 876 A (ESPEY LAWRENCE L) 18 October 1994 (1994-10-18) the whole document	1-45
A	GB 1 065 127 A (SERONO IST FARM) 12 April 1967 (1967-04-12) the whole document	1-45

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP2004/050432

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
EP 0974359	A-	26-01-2000	AT 228850 T	15-12-2002
			AU 4998099 A	14-02-2000
			BR 9912243 A	16-10-2001
			CA 2335340 A1	03-02-2000
			CN 1309567 T	22-08-2001
			DE 69904262 D1	16-01-2003
			DE 69904262 T2	24-07-2003
			DK 974359 T3	03-03-2003
			EP 1188444 A1	20-03-2002
			EP 0974359 A2	26-01-2000
			ES 2183486 T3	16-03-2003
			HK 1026615 A1	24-10-2003
			ID 28747 A	28-06-2001
			JP 2002521342 T	16-07-2002
			NO 20010326 A	20-03-2001
			NZ 508874 A	26-03-2004
			PL 346127 A1	28-01-2002
			PT 974359 T	28-02-2003
			SI 974359 T1	30-04-2003
			WO 0004913 A1	03-02-2000
			US 2003166525 A1	04-09-2003
			US 2002165146 A1	07-11-2002
			ZA 200100190 A	08-01-2002
EP 0853945	A	22-07-1998	EP 0853945 A1	22-07-1998
			EP 1285665 A1	26-02-2003
			AT 237348 T	15-05-2003
			AU 736339 B2	26-07-2001
			AU 5205998 A	23-07-1998
			BR 9800323 A	29-06-1999
			CA 2226803 A1	15-07-1998
			CN 1191139 A	26-08-1998
			CZ 9800135 A3	12-08-1998
			DE 69813322 D1	22-05-2003
			DE 69813322 T2	19-02-2004
			DK 853945 T3	04-08-2003
			ES 2196474 T3	16-12-2003
			HK 1010150 A1	25-07-2003
			HU 9800056 A2	28-08-1998
			ID 19499 A	16-07-1998
			JP 10203997 A	04-08-1998
			NO 980177 A	16-07-1998
			NZ 329572 A	28-05-1999
			PL 324279 A1	20-07-1998
			PT 853945 T	29-08-2003
			RU 2203086 C2	27-04-2003
			TR 9800062 A2	21-08-1998
			TW 518235 B	21-01-2003
			US 5929028 A	27-07-1999
			ZA 9800083 A	08-07-1998
US 5811096	A	22-09-1998	FR 2733914 A1	15-11-1996
			AU 5213996 A	21-11-1996
			CA 2175971 A1	12-11-1996
			CN 1147961 A , B	23-04-1997
			CZ 9601350 A3	11-12-1996
			EA 960024 A2	31-12-1996
			EP 0742013 A1	13-11-1996

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

T/EP2004/050432

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US 5811096	A	GR 3031896 T3 HU 9601241 A2 IL 118171 A JP 2950775 B2 JP 9025242 A NO 961915 A NZ 286554 A PL 314147 A1 SG 42355 A1 AT 183925 T AU 710288 B2 BR 1100337 A3 DE 69604009 D1 DE 69604009 T2 DK 742013 T3 ES 2137635 T3 SI 742013 T1 TW 420614 B ZA 9603683 A	29-02-2000 28-04-1997 31-08-2000 20-09-1999 28-01-1997 12-11-1996 24-09-1998 12-11-1996 15-08-1997 15-09-1999 16-09-1999 25-07-2000 07-10-1999 06-04-2000 06-03-2000 16-12-1999 31-12-1999 01-02-2001 10-11-1997
US 6066620	A 23-05-2000	US 5955427 A US 5696090 A US 5488034 A US 6255284 B1 AT 206308 T AU 695125 B2 AU 6133094 A CA 2155005 A1 CN 1127992 A DE 69428521 D1 DE 69428521 T2 DK 682524 T3 EP 0682524 A1 ES 2164098 T3 HK 1014156 A1 JP 10513433 T PT 682524 T WO 9417819 A1	21-09-1999 09-12-1997 30-01-1996 03-07-2001 15-10-2001 06-08-1998 29-08-1994 18-08-1994 31-07-1996 08-11-2001 23-05-2002 28-01-2002 22-11-1995 16-02-2002 17-04-2003 22-12-1998 28-03-2002 18-08-1994
EP 0318081	A 31-05-1989	AT 65410 T AU 2579188 A CA 1317899 C DE 3863908 D1 DK 648088 A EP 0318081 A1 ES 2051830 T3 GR 3002971 T3 IE 60719 B1 JP 1168624 A MX 165991 B NZ 227088 A US 4902500 A ZA 8808577 A	15-08-1991 01-06-1989 18-05-1993 29-08-1991 28-05-1989 31-05-1989 01-07-1994 25-01-1993 10-08-1994 04-07-1989 14-12-1992 26-07-1990 20-02-1990 30-08-1989
FR 2782455	A 25-02-2000	FR 2782455 A1 AT 230258 T AU 5173699 A DE 69904727 D1	25-02-2000 15-01-2003 14-03-2000 06-02-2003

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

T/EP2004/050432

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
FR 2782455	A		DE 69904727 T2	13-11-2003
			DK 1105102 T3	22-04-2003
			EP 1105102 A1	13-06-2001
			ES 2189456 T3	01-07-2003
			WO 0010534 A1	02-03-2000
			PT 1105102 T	30-04-2003
			SI 1105102 T1	31-08-2003
EP 0603159	A	22-06-1994	US 5096885 A	17-03-1992
			EP 0603159 A2	22-06-1994
			AT 112685 T	15-10-1994
			AU 627174 B2	20-08-1992
			AU 3368789 A	03-11-1989
			CA 1329543 C	17-05-1994
			DE 68918853 D1	17-11-1994
			DE 68918853 T2	13-04-1995
			EP 0409870 A1	30-01-1991
			JP 2747073 B2	06-05-1998
			JP 3503764 T	22-08-1991
			NZ 228747 A	25-09-1991
			WO 8909614 A1	19-10-1989
			US 5763394 A	09-06-1998
			US 2003013653 A1	16-01-2003
			US 2001007858 A1	12-07-2001
EP 0448146	A	25-09-1991	AT 107172 T	15-07-1994
			AU 631730 B2	03-12-1992
			AU 7362391 A	26-09-1991
			CA 2037884 A1	21-09-1991
			DE 69102465 D1	21-07-1994
			DE 69102465 T2	02-03-1995
			DK 448146 T3	24-10-1994
			EP 0448146 A1	25-09-1991
			ES 2057731 T3	16-10-1994
			FI 911309 A	21-09-1991
			HK 1002494 A1	28-08-1998
			ID 905 B	10-09-1996
			IE 910732 A1	25-09-1991
			JP 3031570 B2	10-04-2000
			JP 4217630 A	07-08-1992
			KR 221123 B1	15-09-1999
			NZ 237458 A	25-02-1992
			PT 97074 A , B	31-10-1991
			US 5384132 A	24-01-1995
			US 5270057 A	14-12-1993
			ZA 9101659 A	24-12-1991
US 5650390	A	22-07-1997	IT 1250075 B	30-03-1995
			AT 189608 T	15-02-2000
			AU 677773 B2	08-05-1997
			AU 3266293 A	19-07-1993
			CA 2124801 A1	24-06-1993
			DE 69230672 D1	16-03-2000
			DE 69230672 T2	21-09-2000
			DK 618808 T3	15-05-2000
			EP 0618808 A1	12-10-1994
			ES 2142860 T3	01-05-2000
			GR 3033276 T3	29-09-2000

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

T/EP2004/050432

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
US 5650390	A		WO 9311788 A1	24-06-1993
			JP 7507047 T	03-08-1995
			PT 618808 T	31-07-2000
US 5356876	A	18-10-1994	AU 3423693 A	03-08-1993
			CA 2128017 A1	22-07-1993
			EP 0623027 A1	09-11-1994
			JP 7507050 T	03-08-1995
			WO 9313799 A1	22-07-1993
GB 1065127	A	12-04-1967	FR 3819 M	
			NL 6402059 A	14-09-1964